

# Complete Sequence of a *bla*<sub>NDM-1</sub>-Harboring Plasmid in an *Acinetobacter bereziniae* Clinical Strain Isolated in Argentina

Marco Brovedan,<sup>a</sup> Patricia M. Marchiaro,<sup>a</sup> Jorgelina Morán-Barrio,<sup>a</sup> Marcela Cameranesi,<sup>a</sup> Gabriela Cera,<sup>b</sup> Mariángel Rinaudo,<sup>b</sup> Alejandro M. Viale,<sup>a</sup> Adriana S. Limansky<sup>a</sup>

Instituto de Biología Molecular y Celular de Rosario (IBR, CONICET), Facultad de Ciencias Bioquímicas y Farmacéuticas, Universidad Nacional de Rosario, Rosario, Argentina<sup>a</sup>; Laboratorio de Bacteriología, Facultad de Ciencias Bioquímicas y Farmacéuticas, Universidad Nacional de Rosario, Rosario, Argentina<sup>b</sup>

The New Delhi metallo- $\beta$ -lactamase (NDM-1) was initially identified in clinical isolates of *Escherichia coli* and *Klebsiella pneumoniae* in Sweden from a patient previously hospitalized in India (1). Since then, *bla*<sub>NDM-1</sub> has frequently been reported in *Enterobacteriaceae* and *Acinetobacter* spp., with a fast dissemination in the Indian subcontinent, the Balkan countries, China, and the Middle East (2). NDM-1 producers generally associated with *Enterobacteriaceae* species have also been reported, albeit with much lower frequency, in Latin American countries, including Guatemala, Mexico, Colombia, and Brazil (3). Moreover, production of NDM-1 in this geographic region has also been noted in *Acinetobacter baumannii* in Honduras and Brazil (3, 4) and in *Acinetobacter pittii* in Paraguay and Brazil (5, 6). Here, we report the first case of an NDM-1-producing *Acinetobacter* species in Argentina, an *Acinetobacter bereziniae* clinical isolate. We describe also the complete sequence of a *bla*<sub>NDM-1</sub>-containing plasmid in this strain.

(Part of this work was presented at the 10th International Symposium on the Biology of *Acinetobacter* 2015, Athens, Greece, 3 to 6 June 2015 [7].)

*A. bereziniae* HPC229 was isolated on June 2014 from a blood sample of a 53-year-old female patient that underwent chemotherapy due to leukemia at a hospital located in Rosario, Argentina. The patient was treated with ciprofloxacin plus tigecycline, resulting in clinical and microbiological cure as evaluated by negative blood cultures. The patient was readmitted to the hospital 3 months later with symptoms of severe sepsis and died due to septic shock, with positive blood cultures that grew *Escherichia coli*.

HPC229, originally identified as *Acinetobacter lwoffii* by the Vitek 2 system (bioMérieux), was reclassified by DNA sequence comparison analyses of its 16S rRNA, *gyrB* (8), and *rpoB* (9) genes. The highest identity was found to the corresponding orthologs of the *A. bereziniae* ATCC 17924 type strain (99.9%, 99.7%, and 99.8%, respectively). The antibiotic susceptibility profile of HPC229 was determined by either the Vitek 2 system or the agar dilution method. The interpretation of the obtained MICs based on CLSI breakpoints (10) indicated resistance to  $\beta$ -lactams, including carbapenems (Table 1).

PCR amplification using specific primers for the *bla*<sub>IMP</sub>, *bla*<sub>VIM</sub>, *bla*<sub>SPM</sub>, *bla*<sub>NDM</sub>, *bla*<sub>OXA-23-like</sub>, *bla*<sub>OXA-24/40-like</sub>, and *bla*<sub>OXA-58-like</sub> genes (11–13) and sequencing analysis revealed the presence of *bla*<sub>NDM-1</sub>. Evidence that *bla*<sub>NDM-1</sub> was located in a plasmid was first obtained by plasmid curing (14) in which HPC229 was cultured in 5 ml of LB liquid medium containing 0.2 ml of 10% SDS for 48 h at 40°C. This procedure allowed the isolation of HPC229c, which showed a marked increase in susceptibility to  $\beta$ -lactams (Table 1). Susceptibility to aztreonam as judged by the disk diffusion assay, in contrast, showed no differ-

TABLE 1 Antimicrobial susceptibilities of *A. bereziniae* HPC229 and its plasmid-cured HPC229c derivative<sup>c</sup>

Antimicrobial(s)	MIC ( $\mu$ g/ml)	
	HPC229	HPC229c
Ampicillin-sulbactam <sup>a</sup>	16	$\leq 2$
Ceftazidime <sup>b</sup>	$\geq 256$	8
Cefotaxime <sup>a</sup>	$\geq 64$	8
Cefepime <sup>b</sup>	$\geq 256$	1
Piperacillin-tazobactam <sup>a</sup>	$\geq 128$	8
Imipenem <sup>b</sup>	128	0.125
Meropenem <sup>a</sup>	$\geq 16$	1
Colistin <sup>a</sup>	$\leq 0.5$	$\leq 0.5$
Gentamicin <sup>a</sup>	$\leq 1$	$\leq 1$
Amikacin <sup>a</sup>	$\leq 2$	$\leq 2$
Trimethoprim-sulfamethoxazole <sup>a</sup>	$\leq 20$	$\leq 20$
Ciprofloxacin <sup>a</sup>	0.5	0.5

<sup>a</sup> MICs were determined by the Vitek 2 system.

<sup>b</sup> MICs were determined by the agar dilution method.

<sup>c</sup> HPC229 was resistant to ampicillin-sulbactam, ceftazidime, cefotaxime, cefepime, piperacillin-tazobactam, imipenem, and meropenem, and it was susceptible to colistin, gentamicin, amikacin, trimethoprim-sulfamethoxazole, and ciprofloxacin according to CLSI breakpoints (10).

ences between HPC229 and HPC229c. All conjugation attempts using HPC229 as the donor and *Escherichia coli* DH5 $\alpha$ , *A. baumannii* ATCC 17978, or *Pseudomonas aeruginosa* PAO1 as the recipient (11) were unsuccessful. The genomic relatedness between HPC229 and HPC229c was confirmed using previously described PCR-based procedures (15), and the loss of *bla*<sub>NDM-1</sub> was confirmed by specific PCR analysis.

Analysis of whole-genome sequencing performed with pyrosequencing (454 Genome Sequencer FLX system; Roche Diagnostics) demonstrated that the *bla*<sub>NDM-1</sub> metallo- $\beta$ -lactamase gene was located in HPC229 in a 44,560-bp plasmid, designated pNDM229. This gene was found located inside a composite transposon bracketed by two copies of IS*Aba125* and identical to that first described in *A. baumannii* (16) and designated Tn125 by Poirel et al. (17). The immediate genetic environment of Tn125 in


Accepted manuscript posted online 27 July 2015

Citation Brovedan M, Marchiaro PM, Morán-Barrio J, Cameranesi M, Cera G, Rinaudo M, Viale AM, Limansky AS. 2015. Complete sequence of a *bla*<sub>NDM-1</sub>-harboring plasmid in an *Acinetobacter bereziniae* clinical strain isolated in Argentina. *Antimicrob Agents Chemother* 59:6667–6669. doi:10.1128/AAC.00367-15.

Address correspondence to Adriana S. Limansky, limansky@ibr-conicet.gov.ar.

Copyright © 2015, American Society for Microbiology. All Rights Reserved.

doi:10.1128/AAC.00367-15



**FIG 1** Comparative analysis of *Acinetobacter*  $bla_{NDM-1}$ -harboring plasmids. The linearized genetic maps of pNDM229 from *A. bereziniae* HPC229 (GenBank accession number [KT072713](https://www.ncbi.nlm.nih.gov/nuccore/KT072713)), pNDM-BJ01 from *A. lwoffii* WJ10621 ([NC\\_019268.1](https://www.ncbi.nlm.nih.gov/nuccore/NC_019268.1) [18]), and pNDM-40-1 from *A. bereziniae* CHI-40-1 ([KF702385](https://www.ncbi.nlm.nih.gov/nuccore/KF702385) [19]) are compared. Gray shading indicates more than 99% nucleotide sequence identity. In all plasmids, the *ISAbal4* element is indicated by black vertical lines. A second copy of *ISAbal4* present only in pNDM229 is shown in dark gray and exhibits 94.0% nucleotide identity with the upstream *ISAbal4* element. Partial deletions in conjugative genes in pNDM229 are indicated by radial gray gradients. Arrowheads indicate the direction of transcription, and open reading frames coding hypothetical proteins are indicated by white arrows.

pNDM229 is most similar to that of the conjugative plasmid pNDM-BJ01 (18) present in an *A. lwoffii* clinical strain (Fig. 1). In both cases, the *Tn125* element is located immediately downstream of an *aphA6* gene, which is in turn adjacent to an upstream *ISAbal4* insertion sequence. In the case of pNDM229, however, a second *ISAbal4* copy was found immediately downstream of the *Tn125* element (Fig. 1), thus representing a novel difference from the similar arrangements previously reported (18, 19). This suggests the evolution of a novel composite transposon now bounded by two *ISAbal4* elements with the potential ability to mobilize as a whole.

In summary, we describe here the first case of an NDM-1-producing *A. bereziniae* isolate in Argentina. To our knowledge, only one other *A. bereziniae* NDM-1-producing clinical isolate has been recently reported (CHI-40-1), and the  $bla_{NDM-1}$  gene was located in this strain in a conjugative plasmid (19). However, a number of differences were noted between HPC229 and CHI-40-1 and between the plasmids carried by them (Fig. 1). First, comparison analyses of the corresponding 16S rRNA, *gyrB*, and *rpoB* genes revealed identities of 99.9%, 99.8%, and 99.6%, respectively, indicating no immediate clonal relationship between these two *A. bereziniae* strains. Second, comparison of the corresponding  $bla_{NDM-1}$ -containing platforms revealed two deletions in pNDM-40-1 which were not observed in pNDM229, one of which removed 1,298 bp from the 3' end of *ble* to *dsbC* and the other of which removed 150 bp within *insE* (Fig. 1). Third, the above-noted second copy of *ISAbal4* downstream of *Tn125* was present only in pNDM229. Fourth, unlike with other conjugative plasmids, such as pNDM-40-1 and pNDM-BJ01, some conjugation-related genes, including *virD4*, *trg*, *virB11*, *virB8*, *virB10*, and *virB9*, were either truncated or absent in pNDM229 (Fig. 1), thus explaining conjugation failures.

The origin of *A. bereziniae* HPC229 remains obscure, since no history of overseas travel of the compromised patient or direct contact with foreigners could be identified. The above findings represent a worrisome concern, especially when considering that *Acinetobacter* species other than *A. baumannii* can act as reservoirs of  $bla_{NDM}$  genes and most probably contribute to their spreading among clinically relevant *Enterobacteriaceae* species (6, 19).

**Nucleotide sequence accession numbers.** The 16S rRNA, *gyrB*, and *rpoB* sequences of the HPC229 strain used for comparison with other *Acinetobacter* species have been released in GenBank under the accession numbers [KP765739](https://www.ncbi.nlm.nih.gov/nuccore/KP765739), [KP765740](https://www.ncbi.nlm.nih.gov/nuccore/KP765740),

and [KP765741](https://www.ncbi.nlm.nih.gov/nuccore/KP765741), respectively. The 44,560 bp  $bla_{NDM-1}$ -containing pNDM229 plasmid sequence has been released under accession number [KT072713](https://www.ncbi.nlm.nih.gov/nuccore/KT072713).

#### ACKNOWLEDGMENTS

We thank the anonymous reviewers for their careful reading of our manuscript and their many insightful comments and suggestions for improvement. We are grateful to the Bacteriology Service personnel of the Hospital Provincial del Centenario of Rosario City for kindly providing the HPC229 strain.

P.M.M., G.C., M.R., and A.S.L. are researchers at the National University of Rosario. J.M.-B. and A.M.V. are staff members of CONICET. M.B. and M.C. are fellows of ANPCyT. This work was supported by grants from the Agencia Nacional de Promoción Científica y Tecnológica (ANPCyT, Argentina), the Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), the Ministerio de Salud and the Secretaría de Ciencia, Tecnología e Innovación, Provincia de Santa Fe, and the Secretaría de Salud Pública of the municipality of Rosario.

#### REFERENCES

1. Yong D, Toleman MA, Giske CG, Cho HS, Sundman K, Lee K, Walsh TR. 2009. Characterization of a new metallo-beta-lactamase gene,  $bla_{NDM-1}$ , and a novel erythromycin esterase gene carried on a unique genetic structure in *Klebsiella pneumoniae* sequence type 14 from India. *Antimicrob Agents Chemother* 53:5046–5054. [http://dx.doi.org/10.1128/AAC.00774-09](https://doi.org/10.1128/AAC.00774-09).
2. Dortet L, Poirel L, Nordmann P. 2014. Worldwide dissemination of the NDM-type carbapenemases in Gram-negative bacteria. *Biomed Res Int* 2014:249856. [http://dx.doi.org/10.1155/2014/249856](https://doi.org/10.1155/2014/249856).
3. Pillonetto M, Arend L, Vespero EC, Pelisson M, Chagas TP, Carvalho-Assef AP, Asensi MD. 2014. First report of NDM-1-producing *Acinetobacter baumannii* sequence type 25 in Brazil. *Antimicrob Agents Chemother* 58:7592–7594. [http://dx.doi.org/10.1128/AAC.03444-14](https://doi.org/10.1128/AAC.03444-14).
4. Waterman PE, McGann P, Snesrud E, Robert J, Clifford RJ, Kwak YI, Munoz-Urbizo IP, Tabora-Castellanos J, Milillo M, Preston L, Aviles R, Sutter DE, Leshoa EP. 2013. Bacterial peritonitis due to *Acinetobacter baumannii* sequence type 25 with plasmid-borne New Delhi metallo-beta-lactamase in Honduras. *Antimicrob Agents Chemother* 57:4584–4586. [http://dx.doi.org/10.1128/AAC.00275-13](https://doi.org/10.1128/AAC.00275-13).
5. Pasteran F, Mora MM, Albornoz E, Faccone D, Franco R, Ortellado J, Melgarejo N, Gomez S, Riquelme I, Matheu J, Ramon-Pardo P, Corso A. 2014. Emergence of genetically unrelated NDM-1-producing *Acinetobacter pittii* strains in Paraguay. *J Antimicrob Chemother* 69:2575–2578. [http://dx.doi.org/10.1093/jac/dku139](https://doi.org/10.1093/jac/dku139).
6. Pagano M, Poirel L, Martins AF, Rozales FP, Zavascki AP, Barth AL, Nordmann P. April 2015. Emergence of NDM-1-producing *Acinetobacter pittii* in Brazil. *Int J Antimicrob Agents* [http://dx.doi.org/10.1016/j.ijantimicag.2014.12.011](https://doi.org/10.1016/j.ijantimicag.2014.12.011).
7. Brovedan M, Marchiaro PM, Morán-Barrio J, Cameranesi M, Cera G,

- Rinaudo M, Viale AM, Limansky AS. 2015. 10th International Symposium on the Biology of *Acinetobacter* 2015, Athens, Greece, 3 to 6 June 2015, abstr P33A.
8. Ibrahim A, Gerner-Smidt P, Liesack W. 1997. Phylogenetic relationship of the twenty-one DNA groups of the genus *Acinetobacter* as revealed by 16S ribosomal DNA sequence analysis. *Int J Syst Bacteriol* 47:837–841. <http://dx.doi.org/10.1099/00207713-47-3-837>.
  9. Bonnin RA, Ocampo-Sosa AA, Poirer L, Guet-Revillet H, Nordmann P. 2012. Biochemical and genetic characterization of carbapenem-hydrolyzing  $\beta$ -lactamase OXA-229 from *Acinetobacter bereziniae*. *Antimicrob Agents Chemother* 56:3923–3927. <http://dx.doi.org/10.1128/AAC.00257-12>.
  10. Clinical and Laboratory Standards Institute (CLSI). 2014. Performance standards for antimicrobial susceptibility testing. Twenty-third informational supplement. Document M100-S24. CLSI, Wayne, PA.
  11. Marchiaro P, Viale AM, Ballerini V, Rossignol G, Vila AJ, Limansky A. 2010. First report of a Tn402-like class 1 integron carrying *bla*<sub>VIM-2</sub> in *Pseudomonas putida* from Argentina. *J Infect Dev Ctries* 4:412–416.
  12. Manchanda V, Rai S, Gupta S, Rautela RS, Chopra R, Rawat DS, Verma N, Singh NP, Kaur IR, Bhalla P. 2011. Development of TaqMan real-time polymerase chain reaction for the detection of the newly emerging form of carbapenem resistance gene in clinical isolates of *Escherichia coli*, *Klebsiella pneumoniae*, and *Acinetobacter baumannii*. *Indian J Med Microbiol* 29:249–253. <http://dx.doi.org/10.4103/0255-0857.83907>.
  13. Marchiaro P, Ballerini V, Spalding T, Cera G, Mussi MA, Morán-Barrío J, Vila AJ, Viale AM, Limansky AS. 2008. A convenient microbiological assay employing cell-free extracts for the rapid characterization of Gram-negative carbapenemase producers. *J Antimicrob Chemother* 62:336–344. <http://dx.doi.org/10.1093/jac/dkn185>.
  14. Lopes BS, Al-Hassan L, Amyes SG. 2012. *ISAbas825* controls the expression of the chromosomal *bla*<sub>OXA-51-like</sub> and the plasmid borne *bla*<sub>OXA-58</sub> gene in clinical isolates of *Acinetobacter baumannii* isolated from the USA. *Clin Microbiol Infect* 18:E446–E451. <http://dx.doi.org/10.1111/j.1469-0691.2012.03979.x>.
  15. Limansky AS, Viale AM. 2002. Can composition and structural features of oligonucleotides contribute to their wide-scale applicability as random PCR primers in mapping bacterial genome diversity? *J Microbiol Methods* 50:291–297. [http://dx.doi.org/10.1016/S0167-7012\(02\)00040-4](http://dx.doi.org/10.1016/S0167-7012(02)00040-4).
  16. Pfeifer Y, Wilharm G, Zander E, Wichelhaus TA, Göttig S, Hunfeld KP, Seifert H, Witte W, Higgins PG. 2011. Molecular characterization of *bla*<sub>NDM-1</sub> in an *Acinetobacter baumannii* strain isolated in Germany in 2007. *J Antimicrob Chemother* 66:1998–2001. <http://dx.doi.org/10.1093/jac/dkr256>.
  17. Poirer L, Bonnin RA, Boulanger A, Schrenzel J, Kaase M, Nordmann P. 2012. Tn125-related acquisition of *bla*<sub>NDM-like</sub> genes in *Acinetobacter baumannii*. *Antimicrob Agents Chemother* 56:1087–1089. <http://dx.doi.org/10.1128/AAC.05620-11>.
  18. Hu H, Hu Y, Pan Y, Liang H, Wang H, Wang X, Hao Q, Yang X, Yang X, Xiao X, Luan Ch, Yang Y, Cui Y, Yang R, Gao G, Song Y, Zhu B. 2012. Novel plasmid and its variant harboring both a *bla*<sub>NDM-1</sub> gene and type IV secretion system in clinical isolates of *Acinetobacter lwoffii*. *Antimicrob Agents Chemother* 56:1698–1702. <http://dx.doi.org/10.1128/AAC.06199-11>.
  19. Jones LS, Carvalho MJ, Toleman MA, White PL, Connor TR, Mushtaq A, Weeks JL, Kumarasamy KK, Raven KE, Török ME, Peacock SJ, Howe RA, Walsh TR. 2015. Characterization of plasmids in extensively drug-resistant *Acinetobacter* strains isolated in India and Pakistan. *Antimicrob Agents Chemother* 59:923–929. <http://dx.doi.org/10.1128/AAC.03242-14>.