

Draft Genome Sequence of *Lactococcus lactis* subsp. *lactis* bv. *diacetylactis* CRL264, a Citrate-Fermenting Strain

Federico Zuljan,^{a,c} Martín Espariz,^{a,c} Victor S. Blancato,^{a,c} Luis Esteban,^{c,d} Sergio Alarcón,^{b,c} Christian Magni^{a,c}

Laboratorio de Fisiología y Genética de Bacterias Lácticas, Instituto de Biología Molecular y Celular de Rosario (IBR-CONICET), Rosario, Argentina^a; Instituto de Química de Rosario (QUIR-CONICET), Rosario, Argentina^b; Laboratorio Biotecnología e Inocuidad de los Alimentos, Facultad de Ciencias Bioquímicas y Farmacéuticas, Universidad Nacional de Rosario (UNR), Rosario, Argentina^c; Facultad de Ciencias Médicas, UNR, Rosario, Argentina^d

We report the draft genome sequence of *Lactococcus lactis* subsp. *lactis* bv. *diacetylactis* CRL264, a natural strain isolated from artisanal cheese from northwest Argentina. *L. lactis* subsp. *lactis* bv. *diacetylactis* is one of the most important microorganisms used as starter culture around the world. The CRL264 strain constitutes a model microorganism in the studies on the generation of aroma compounds (diacetyl, acetoin, and 2,3-butanediol) by lactic acid bacteria. Our genome analysis shows similar genetic organization to other available genomes of *L. lactis* bv. *diacetylactis* strains.

Received 18 November 2015 Accepted 9 December 2015 Published 4 February 2016

Citation Zuljan F, Espariz M, Blancato VS, Esteban L, Alarcón S, Magni C. 2016. Draft genome sequence of *Lactococcus lactis* subsp. *lactis* bv. *diacetylactis* CRL264, a citrate-fermenting strain. *Genome Announc* 4(1):e01575-15. doi:10.1128/genomeA.01575-15.

Copyright © 2016 Zuljan et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Christian Magni, magni@ibr-conicet.gov.ar.

Lactococcus lactis is a mesophilic bacterium widely used in the production of fermented milk products, particularly cheese. Technological strains are divided into three phenotypes: *L. lactis*, *L. cremoris*, and *L. diacetylactis*. *L. lactis* IL1403 is a plasmid-free strain derived from *L. lactis* subsp. *lactis* bv. *diacetylactis* IL594 (*L. diacetylactis*) (1) and constitutes a model microorganism, being the first sequenced genome of a lactic acid bacterium (2).

The *L. diacetylactis* CRL264 strain was initially characterized for its capacity to ferment citrate (3), and it was later extensively characterized in our labs at the biochemical and genetic levels (4–11). Citrate is transported by a plasmid-encoded transporter (CitP) (4–6), whereas genes encoding the enzymes required for the intracellular metabolism of citrate are located in a large chromosomal operon (8). The citrate lyase enzyme complex is responsible for cleaving citrate into acetate and oxaloacetate, which is subsequently decarboxylated to pyruvate by the action of the soluble oxaloacetate decarboxylase (7, 9). Also, it has been demonstrated that acidification of the external medium is required for the induction of both transporter (plasmidic) and catabolic genes (chromosomal) (6–8, 10). Moreover, the route involved in the generation of the aroma compounds (*butBA* encoding 2,3-butanediol dehydrogenase and diacetyl-acetoin reductase, *als* encoding acetolactate synthetase, and *aldB* and *aldC* both encoding acetolactate decarboxylase) is also induced at low pH (10, 11).

In this report, the draft genome sequence of *L. diacetylactis* CRL264, consisting of 83 contigs, is presented. The genome sequence was determined by using an Illumina HiSeq 2000 platform (MR DNA, USA). Base-calling was carried out with HiSeq control software version 1.4.8. (Metrics sum, 2.60 Mbp; N_{50} , 48.2 kbp; max, 252.7 kbp; min, 200 bp; G+C content, 36.4%; number of clusters, 3,891; number of reads, 7,782; size: 463.57 Mb; coverage, 424.68×. The short reads were *de novo* assembled with SeqMan NGen (DNASTAR Inc.). BLASTn analysis (all versus all) with the resulting contigs was performed, and those shorter than 1,000 bp and with an identity higher than 99% with sequences already con-

tained in a longer contig were deleted. The remaining contigs were ordered and oriented with Advanced Pipemaker (12) and Mauve version 2.3.1 (13) using the annotated genome of *L. lactis* IL1403 as the genome of reference for *L. diacetylactis* CRL264.

The complete genome of strain CRL264 contains a total of 2,500 coding sequences; 63 structural RNAs were detected, and over 45% of the genes were assigned to specific subsystem categories by RAST (Rapid Annotations using Subsystem Technology) (14) and BASys (Bacterial Annotation System) (15). Manual curation of the genes was performed with the Seed viewer version 2.0, comparing the presence and absence of genes with closely related species. Comparative analysis of the chromosomal genome of the CRL264 strain among *L. diacetylactis* strains IL1403, LD61, TIFN2, and TIFN4 showed a clonal origin of the diacetylactis strain. Experimental and genomic comparative analysis will provide further insight into the strain's capacity to produce diacetyl.

Nucleotide sequence accession number. The complete genome of *L. lactis* bv. *diacetylactis* CRL264 has been deposited at DDBJ/EMBL/GenBank under accession number [LKPE00000000](https://www.ncbi.nlm.nih.gov/nuccore/LKPE00000000).

ACKNOWLEDGMENTS

This work was supported by grants from the Agencia Nacional de Promoción Científica y Tecnológica (ANPCyT and PICT 2014-1513 and PICT 2014-3482, Argentina) and CONICET (PIP 2012-2014). F.Z. is a fellow of CONICET; M.E., V.S.B., S.A., and C.M. are researchers at the same institution.

FUNDING INFORMATION

MINCYT | Agencia Nacional de Promoción Científica y Tecnológica (ANPCyT) provided funding to Christian Magni under grant number PICT 2014-1513. MINCYT | Agencia Nacional de Promoción Científica y Tecnológica (ANPCyT) provided funding to Victor Blancato under grant number PICT 2014-3482.

REFERENCES

- Górecki RK, Koryszewska-Bagińska A, Gołębiewski M, Żylińska J, Grynberg M, Bardowski JK. 2011. Adaptive potential of the *Lactococcus*

- lactis* IL594 strain encoded in its 7 plasmids. PLoS One 6: <http://dx.doi.org/10.1371/journal.pone.0022238>.
2. Bolotin A, Wincker P, Mauer S, Jaillon O, Malarme K, Weissenbach J, Ehrlich SD, Sorokin A. 2001. The complete genome sequence of the lactic acid bacterium *Lactococcus lactis* ssp. *lactis* IL1403. Genome Res 11: 731–753.
 3. Sesma F, Gardiol D, de Ruiz Holgado AP, de Mendoza D. 1990. Cloning of the citrate permease gene of *Lactococcus lactis* subsp. *lactis* biovar diacetylactis and expression in *Escherichia coli*. Appl Environ Microbiol 56: 2099–2103.
 4. Magni C, de Felipe FL, Sesma F, López P, de Mendoza D. 1994. Citrate transport in *Lactococcus lactis* biovar diacetylactis: expression of the plasmid-borne citrate permease P. FEMS Microbiol Lett 118:75–82. <http://dx.doi.org/10.1111/j.1574-6968.1994.tb06806.x>.
 5. de Felipe FL, López P, Magni C, de Mendoza D. 1996. Transcriptional activation of the citrate permease P gene of *Lactococcus lactis* biovar diacetylactis by an insertion sequence-like element present in plasmid pCIT264. Molec Gen Genet 250:428–436. <http://dx.doi.org/10.1007/BF02174031>.
 6. García-Quintáns N, Magni C, de Mendoza D, López P. 1998. The citrate transport system of *Lactococcus lactis* subsp. *lactis* biovar diacetylactis is induced by acid stress. Appl Environ Microbiol 64:850–857.
 7. Magni C, de Mendoza D, Konings WN, Lolkema JS. 1999. Mechanism of citrate metabolism in *Lactococcus lactis*: resistance against lactate toxicity at low pH. J Bacteriol 181:1451–1457.
 8. Martín MG, Sender PD, Peirú S, de Mendoza D, Magni C. 2004. Acid-inducible transcription of the operon encoding the citrate lyase complex of *Lactococcus lactis* biovar diacetylactis CRL264. J Bacteriol 186: 5649–5660. <http://dx.doi.org/10.1128/JB.186.17.5649-5660.2004>.
 9. Sender PD, Martín MG, Peirú S, Magni C. 2004. Characterization of an oxaloacetate decarboxylase that belongs to the malic enzyme family. FEBS Lett 570:217–222. <http://dx.doi.org/10.1016/j.febslet.2004.06.038>.
 10. García-Quintáns N, Repizo G, Martín M, Magni C, López P. 2008. Activation of the diacetyl/acetoin pathway in *Lactococcus lactis* subsp. *lactis* bv. diacetylactis CRL264 by acidic growth. Appl Environ Microbiol 74: 1988–1996. <http://dx.doi.org/10.1128/AEM.01851-07>.
 11. Zuljan FA, Repizo GD, Alarcon SH, Magni C. 2014. α -Acetolactate synthase of *Lactococcus lactis* contributes to pH homeostasis in acid stress conditions. Int J Food Microbiol 188:99–107. <http://dx.doi.org/10.1016/j.ijfoodmicro.2014.07.017>.
 12. Schwartz S, Zhang Z, Frazer KA, Smit A, Riemer C, Bouck J, Gibbs R, Hardison R, Miller W. 2000. PipMaker—a Web server for aligning two genomic DNA sequences. Genome Res 10:577–586. <http://dx.doi.org/10.1101/gr.10.4.577>.
 13. Darling AE, Mau B, Perna NT. 2010. progressiveMauve: multiple genome alignment with gene gain, loss and rearrangement. PLoS One 5: <http://dx.doi.org/10.1371/journal.pone.0011147>.
 14. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: Rapid Annotations using Subsystems Technology. BMC Genomics 9: <http://dx.doi.org/10.1186/1471-2164-9-75>.
 15. Van Domselaar GH, Stothard P, Shrivastava S, Cruz JA, Guo A, Dong X, Lu P, Szafron D, Greiner R, Wishart DS. 2005. BASys: a web server for automated bacterial genome annotation. Nucleic Acids Res 33: W455–W459. <http://dx.doi.org/10.1093/nar/gki593>.