


Accepted Manuscript

Title: “The heme uptake process in *Trypanosoma cruzi* epimastigotes is inhibited by heme analogues and by inhibitors of ABC transporters”

Authors: Mauricio Cupello Peixoto, Cíntia Fernandes de Souza, Celeste Buchensky, Juliana Baptista Rocha Corrêa Soares, Gustavo Augusto Travassos Laranja, Marsen Garcia Pinto Coelho, Julia A. Cricco, Marcia Cristina Paes



PII: S0001-706X(11)00253-1
DOI: doi:10.1016/j.actatropica.2011.08.011
Reference: ACTROP 2723

To appear in: *Acta Tropica*

Received date: 27-4-2011
Revised date: 18-7-2011
Accepted date: 20-8-2011

Please cite this article as: Peixoto, M.C., Souza, C.F., Buchensky, C., Soares, J.B.R.C., Laranja, G.A.T., Coelho, M.G.P., Cricco, J.A., Paes, M.C., “The heme uptake process in *Trypanosoma cruzi* epimastigotes is inhibited by heme analogues and by inhibitors of ABC transporters”, *Acta Tropica* (2010), doi:10.1016/j.actatropica.2011.08.011

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.


The heme uptake process in *Trypanosoma cruzi* epimastigotes is inhibited by heme analogues and by inhibitors of ABC transporters Mauricio Cupello Peixoto¹, Cíntia Fernandes de Souza¹, Celeste Buchensky², Juliana Baptista Rocha Corrêa Soares³, Gustavo Augusto Travassos Laranja¹, Marsen Garcia Pinto Coelho¹, Julia A. Cricco², Marcia Cristina Paes^{1,4}

¹Laboratório de Interação Tripanossomatídeos e Vetores - Departamento de Bioquímica, Instituto de Biologia Roberto Alcântara Gomes - IBRAG – Universidade do Estado do Rio de Janeiro – Rio de Janeiro, Brasil; ²Departamento de Química Biológica and Instituto de Biología Molecular y Celular de Rosario (IBR, CONICET-UNR), Facultad de Ciencias Bioquímicas y Farmacéuticas, Universidad Nacional de Rosario, Rosario, Argentina. ³Laboratório de Bioquímica Redox, Programa de Biología Molecular e Biotecnologia, Instituto de Bioquímica Médica, Universidade Federal do Rio de Janeiro, Rio de Janeiro, RJ, Brasil; ⁴ Instituto Nacional de Ciência e Tecnologia - Entomologia Molecular (INCT-EM) - Brasil

Address correspondence to: Marcia Cristina Paes, Boulevard 28 de setembro, 87 fundos IBRAG 4^o andar Bioquímica, Rio de Janeiro, Brasil, Tel.: +55-21-25876143; Fax: +55-21-25876136, E-mail address: mcpaes@uerj.br

Julia A. Cricco, Departamento de Química Biológica and Instituto de Biología Molecular y Celular de Rosario (IBR, CONICET-UNR), Facultad de Ciencias Bioquímicas y Farmacéuticas, Universidad Nacional de Rosario, Suipacha 531, S2002LRK Rosario, Argentina. Tel.: 154 341 435 1235, ext. 109; fax: 154 341 439 0465; e-mail: cricco@ibr.gov.ar

Pictogram:



This is the first indication of the inhibition of the heme transport by analogues in *T. cruzi*, possibly an ABC transporter, which can interfere with the vector – parasite interaction.

< A common porphyrin transporter might be involved in the entrance of amphipathic porphyrins.

< The heme transport blockage in *T. cruzi* epimastigotes affects its biological cycle since decreases the epimastigotes growth.

<The heme uptake process is affected by the presence of different transporter inhibitors.

< Evidences indicate that heme translocation may be mediated by an ABC transporter.

Accepted Manuscript

1 “The heme uptake process in *Trypanosoma cruzi* epimastigotes is inhibited by
2 heme analogues and by inhibitors of ABC transporters”
3
4
5

6 Mauricio Cupello Peixoto¹, Cíntia Fernandes de Souza¹, Celeste Buchensky², Juliana Baptista
7 Rocha Corrêa Soares³, Gustavo Augusto Travassos Laranja¹, Marsen Garcia Pinto Coelho¹,
8
9 Julia A. Cricco^{2j}, and Marcia Cristina Paes^{1,4j}
10
11
12
13
14
15

16 ¹Laboratório de Interação Tripanossomatídeos e Vetores, Departamento de Bioquímica,
17 Instituto de Biologia Roberto Alcântara Gomes, IBRAG, Universidade do Estado do
18 Rio de Janeiro, Rio de Janeiro, Brasil; ² Departamento de Química Biológica and
19 Instituto de Biología Molecular y Celular de Rosario (IBR, CONICET-UNR), Facultad
20 de Ciencias Bioquímicas y Farmacéuticas, Universidad Nacional de Rosario, Rosario,
21 Argentina.³ Laboratório de Bioquímica Redox, Programa de Biologia Molecular e
22 Biotecnologia, Instituto de Bioquímica Médica, Universidade Federal do Rio de Janeiro,
23 Rio de Janeiro, RJ, Brasil; ⁴Instituto Nacional de Ciência e Tecnologia - Entomologia
24 Molecular (INCT-EM) – Brasil.
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42

43 ^jAddress correspondence to: Marcia Cristina Paes, Boulevard 28 de setembro, 87 fundos
44 IBRAG 4^o andar Bioquímica, Rio de Janeiro, Brasil, Tel.: +55-21-25876143; Fax: +55-21-
45 25876136, E-mail address: mcpaes@uerj.br
46
47
48
49
50

51 Julia A. Cricco, Departamento de Química Biológica and Instituto de Biología
52 Molecular y Celular de Rosario (IBR, CONICET-UNR), Facultad de Ciencias
53 Bioquímicas y Farmacéuticas, Universidad Nacional de Rosario, Suipacha 531,
54
55
56
57
58
59
60
61
62
63
64
65

S2002LRK Rosario, Argentina. Tel.: 154 341 435 1235, ext. 109; Fax: 154 341 439

0465; E-mail: crizzo@ibr.gov.ar

Accepted Manuscript

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Abstract

1
2
3 Heme (iron protoporphyrin IX) is an important molecule involved in many biological
4
5 reactions, including oxygen transport, respiration, photosynthesis and drug detoxification.
6
7
8 *Trypanosoma cruzi* parasites, the etiological agent of Chagas' disease, take up heme from
9
10 the environment to supply their nutritional needs because they do not synthesize this
11
12 cofactor. However, the mechanisms involved in heme transport across biological
13
14 membranes are poorly understood. Indeed, in *T. cruzi*, no heme transporter has yet been
15
16 characterized. In the present work, we evaluate the heme uptake processes by *T. cruzi*
17
18 epimastigotes using fluorescent heme-analogues. Heme uptake decreased significantly when
19
20 cells were pretreated with different concentrations of SnPPIX, PdMPIX or ZnMPIX, this
21
22 observed competition suggests that they are taken up by the same transport system. We
23
24 studied the growth behavior of epimastigotes using the same heme-analogues and the
25
26 treatments with SnPPIX or PdMPIX impaired cell growth but when heme was added to the
27
28 culture medium the observed inhibition was partially reversed. In addition, we tested how
29
30 the heme uptake processes are affected by the presence of different transporter inhibitors.
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

When the cells were treated with inhibitors and then incubated with heme, heme uptake decreased significantly for all treatments. These results constitute a strong indication for the existence of a protein associated with porphyrin transport in *T. cruzi*, possibly ATP-binding cassette transporters (ABC-transporter).

Keywords:

Heme; *Trypanosoma cruzi*; porphyrins; heme transport

Introduction

1
2
3 *Trypanosoma cruzi* is a protozoan parasite that resides in the blood and other tissues
4
5 and causes American Trypanosomiasis, or Chagas' disease, which is recognized by the World
6
7 Health Organization (WHO) as one of thirteen major neglected diseases throughout the world.
8
9 Chagas' disease is a significant problem in Latin America (Hotez *et al.*, 2007), where it is
10
11 endemic, but other countries (considered non-endemic) are significantly affected as well
12
13 because of the migration movements of the Latin American population (Coura and Viñas,
14
15 2010). Currently, an estimated 10 million people are infected worldwide, primarily in Latin
16
17 America, and more than 25 million people are at risk of being infected (WHO, 2010). The
18
19 treatment is not very effective; the available medications have significant side effects and little
20
21 efficacy during the chronic stage of the disease. As there are currently no prospects for vaccines
22
23 or satisfactory medical treatments, the search for new therapies is a priority. This disease is
24
25 transmitted by triatomine vector insects while they feed on a vertebrate host. These
26
27 bloodsucking insects ingest 6 to 12 times their original weight in blood (Wigglesworth, 1943).
28
29 This blood is continuously digested through a variety of proteinases, releasing amino acids,
30
31 peptides and heme. Usually, approximately 10 mM of heme bound to hemoglobin is obtained
32
33 in a single ingest (Graça-Souza *et al.*, 2006).
34
35
36
37
38
39
40
41

42 The heme molecule is an important cofactor that is involved in several essential
43
44 biological processes in aerobic organisms, such as oxygen transport (hemoglobin and
45
46 myoglobin), cellular respiration (cytochrome *c*, cytochrome *c* oxidase, and cytochrome *c*
47
48 reductase), drug detoxification (enzyme P450) and enzymes involved in antioxidant defense
49
50 (catalases and peroxidases) (Ponka, 1999). Heme is a highly toxic molecule due to its pro-
51
52 oxidant properties (Ryter and Tyrrel, 2000; Deterding *et al.*, 2004), its biosynthesis and
53
54 degradation are tightly regulated in accordance with cellular requirements. The heme
55
56 biosynthetic pathway is conserved throughout evolution, in non-photosynthetic eukaryotic
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

cells, it starts and finishes in the mitochondria, while still involving several cytosolic steps (Severance and Hamza, 2009). There are some organisms that depend on essential heme-proteins but lack in part or in total a heme biosynthetic pathway. Trypanosomatids are included in this group (revisited by Koreny, 2010; Tripodi *et al.*, 2011). Biochemical studies have demonstrated the absence of a complete heme biosynthesis pathway in *T. cruzi* (Lombardo *et al.*, 2003), and this was later corroborated by the absence of conserved enzymes in its genomic sequence (El-Sayed *et al.*, 2005).

As heme is an essential cofactor for trypanosomatids, the processes of heme binding, transport and intracellular distribution must involve critical pathways in these organisms. Heme uptake in trypanosomatids is a topic of actual debate and, specifically for *T. cruzi*, there is limited information concerning transporters or carriers and there is no biochemical characterization of the process itself. Our group has demonstrated that in *T. cruzi* epimastigotes, heme and hemoglobin internalization proceed through different routes and/or mechanisms (Lara *et al.*, 2007). We have observed by fluorescence microscopy using fluorescent heme analogues that heme uptake might involve the activity of a P-glycoprotein (Pgp) homologue, an ABC transporter. Latter, the protein TbHpHbR was described as a receptor for the complex haptohemoglobin-hemoglobin (HpHb) (Vanhollebeke *et al.*, 2008). The authors demonstrated that TbHpHbR is a bloodstream stage-specific protein, but it is not expressed in the procyclic form. TbHpHbR is present in *T. brucei subsp. gambiense* and *T. brucei subsp. rhodesiense* but absent from the related kinetoplastids *T. cruzi* and *Leishmania*. This protein directs the internalization of heme carried by the Hp-Hb complex into hemoproteins in order to optimize growth of bloodstream forms.

Recently, it was demonstrated that an ATP-binding cassette protein was involved in intracellular heme trafficking in *Leishmania* (Campos-Salinas *et al.*, 2011). The protein LABCG5 was associated with the internal distribution of hemoglobin-bound heme to the

1 mitochondria, but LABCG5 was not involved in the internal trafficking of free heme.
2
3 Currently, the mechanisms by which heme is transported through the plasma membrane and
4
5 targeted to heme proteins in the intracellular medium remain unknown for all the
6
7 trypanosomatids.
8

9
10 Considering the implications of new discoveries in the transport of porphyrins in *T.*
11
12 *cruzi*, it becomes of great importance, especially from the standpoint of chemotherapy, to
13
14 elucidate the mechanisms for heme transport. In this study, we investigate the effects of heme
15
16 transport blockage by heme analogues and specific transporters inhibitors and, consequently,
17
18 the disruption of the life-cycle of *T. cruzi*.
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Materials and methods

1
2 **2.1. Reagents.** BHI (brain-heart infusion medium) was obtained from DIFCO (Sparks, MD,
3 USA). cyclosporin A (CsA) was purchased from Novartis Pharma S.A.S. (Huningue, France).
4 Fetal calf serum (FCS) was purchased from CULTLAB (São Paulo, Brazil). Heme (Fe-
5 protoporphyrin IX), Sn-protoporphyrin IX (SnPPIX), Pd-mesoporphyrin IX (PdMPIX) and Zn-
6 mesoporphyrin IX (ZnMPIX) were obtained from Frontier Scientific (Logan, UT, USA).
7 Indomethacin (1-[p-Chlorobenzoyl]-5-methoxy-2-methylindole-3-acetic acid) was obtained
8 from SIGMA-ALDRICH CHEME (Steinheim, Germany). Pyridine was purchased from
9 MERCK (Darmstadt, Germany). verapamil (5-[N-(3,4-Dimethoxyphenylethyl)methylamino]-
10 2-(3,4-dimethoxyphenyl)-2-isopropylvaleronitrile hydrochloride) was also purchased from
11 SIGMA-ALDRICH CHEME (Steinheim, Germany). All other reagents used were of analytical
12 grade.

13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31 **2.2. Parasites.** *Trypanosoma cruzi* epimastigotes, strain Dm28c, were grown at 28°C for seven
32 days in BHI supplemented with 10% FCS and in the absence or presence of 30 µM heme. The
33 growth was monitored by cell counting in a Neubauer chamber.

34
35
36
37
38
39
40
41 **2.3. Porphyrins.** Fe-protoporphyrin IX stock (20 mM) was prepared by dissolving in 0.1 N
42 NaOH, and after it was buffered using PBS (100 mM sodium phosphate buffer and 150 mM
43 NaCl at pH 7.4). The stock was diluted immediately before use to 10 mM in the same PBS
44 buffer. Stocks of Sn-protoporphyrin IX, Pd-mesoporphyrin IX and Zn-mesoporphyrin IX
45 (heme analogues) were prepared in 0.1 N NaOH and buffered in PBS in the same way as the
46 heme and then used in the experiments.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

2.4. ABC transporter inhibitors (cyclosporin A, indomethacin and verapamil). The inhibitor stock solutions were prepared as follows: the 5 mM cyclosporin A and 10 mM verapamil stock solutions were prepared in water, and the 10 mM indomethacin stock solution was prepared in 0.1 N NaOH and PBS.

2.5. The effect of SnPPIX, PdMPIX, and ZnMPIX on *T. cruzi* epimastigote heme uptake.

The parasites were maintained in BHI supplemented with 10% FCS for 7 days. After 7 days, the cells were washed twice with PBS. After washing, the cells were suspended with PBS and incubated for 10 minutes with different concentrations of the heme analogues (SnPPIX, PdMPIX, and ZnMPIX). After incubation, they were washed twice with PBS, suspended with PBS and then incubated with 10 μ M heme for 10 min. After this incubation, the cells were collected, washed twice with PBS and lysed through repeated freezing and thawing. Heme uptake was measured in epimastigote lysates with the alkaline pyridine method (Falk, 1964) using a GBC UV/VIS 920 absorption spectrometer with the following volume modifications: 500 μ L of sample was combined in a 1 mL cuvette with 500 μ L of stock solution containing 0.2 N NaOH and 48% pyridine. The oxidized and reduced (via the addition of DTH) spectra were recorded between 500 nm and 600 nm. The 557 and 541 nm peaks were indentified in the differential reduced minus oxidized spectra. The heme concentration was estimated using the molar extinction coefficient 20.7 $\text{mM}^{-1} \text{cm}^{-1}$. As controls we recorded the reduced minus oxidized spectra of heme alone and heme plus different amounts of SnPPIX and we did not observe any difference in the reduced minus oxidized spectra, indicating that these analogues did not show interference in this test.

2.6. Analysis of ZnMPIX uptake by *T. cruzi* epimastigotes measured via direct fluorescence from total cell extracts.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

The *T. cruzi* epimastigotes were maintained in BHI supplemented with 10% FCS and 30 μM hemin at 28°C. Parasites in the stationary phase were used to study the ZnMPIX-uptake processes. The cells were collected by centrifugation and washed twice with PBS. The cells were then suspended in PBS, and the assays were performed in PBS buffer at 28°C using 2×10^7 parasites per assay, volume of 100 μl . The epimastigotes were incubated for 5 minutes with 10 μM ZnMPIX at different concentrations of hemin (0–100 μM) added to epismatigotes to complete a volume of 200 μl . After incubation, the reaction was stopped by the addition of 800 μl of cold PBS with 1mM hemin. Then the cells were collected and washed 3 times with cold PBS (we tested that the last washing buffer does not contain hemin nor ZnMPIX). The cells were then suspended in lysis buffer (0.5% Triton X-100 in PBS) and disrupted by repeated freezing and thawing. The lysates were clarified by centrifugation, and the supernatant was used to measure the ZnMPIX fluorescence intensity in the total cell extracts using a Varian Eclipse fluorometer ($\lambda_{\text{ex}} = 405 \text{ nm}$ and $\lambda_{\text{em}} = 578 \text{ nm}$). The fluorescence intensity measured is directly proportional to the ZnMPIX concentration. Every data point is the average of 3 independent assays, and the uptake experiments were reproduced at least 3 times. The same experiments were carried out in stationary phase and we observed the same effects on ZnMPIX uptake.

2.7. The effects of SnPPIX, PdMPIX, and ZnMPIX on *T. cruzi* epimastigote growth.

Parasites were maintained at 28°C for a week in BHI containing both 10% FCS and 30 μM heme, and then epimastigotes in the stationary phase were collected by centrifugation and washed three times with BHI without FCS. The cells were suspended in BHI with 10% FCS. The heme analogues (SnPPIX, PdMPIX, and ZnMPIX) were added to cells suspensions at different concentrations and were incubated for 10 min. Then 30 μM heme was added to the

1 medium. The cells were then maintained for 12 days in culture. Cell growth was monitored by
2 cell counting in a Neubauer chamber.
3

4
5
6
7 **2.8. The effect of cyclosporin A, indomethacin and verapamil on epimastigotes heme**
8 **uptake.** Parasites were maintained in BHI supplemented with 10% FCS. The cells were
9 washed twice with PBS and then treated with different ABC-transporter modulators (300 μ M
10 indomethacin, 10 μ M cyclosporin A or 2 μ M verapamil) for 30 min. Then 10 μ M heme was
11 added to the medium, and the samples were incubated for another 10 min. After this incubation,
12 the cells were lysed through a freeze/thaw process. The heme uptake was measured from the *T.*
13 *cruzi* lysates as previously described in 2.5.
14
15
16
17
18
19
20
21
22
23
24
25
26

27 **2.9. Evaluation of the effect of cyclosporin A on the proliferation of *T. cruzi* epimastigotes.**

28 The cells were maintained at 28° for a week in BHI containing both 10% FCS and 30 μ M
29 heme. During the stationary phase, they were collected by centrifugation and washed three
30 times with BHI without FCS. The cells were suspended in BHI with 10% of FCS. The
31 epimastigotes were pre-treated with increasing doses of cyclosporin A (1 μ M to 20 μ M) for 30
32 minutes, and then 10 μ M heme was added to the cells. The cells were maintained for 7 days in
33 this culture. Cell growth was monitored by cell counting in a Neubauer chamber.
34
35
36
37
38
39
40
41
42
43
44
45

46 **2.10. Evaluation of the effect of cyclosporin A on the heme uptake in *T. cruzi***

47 **epimastigotes.** Parasites were maintained in BHI supplemented with 10% FCS and 30 μ M
48 heme at 28° for one week. Then, the cells in the stationary phase were collected by
49 centrifugation and washed three times with BHI without FCS. After washing, the epimastigotes
50 were suspended in BHI with 10% FCS and pre-treated with 10 μ M of cyclosporin A for 30
51 minutes. After 30 minutes, 300 μ M heme was added to the medium, and the cells were
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 maintained for 7 days in culture. After 7 days, the cells were collected, washed twice with PBS,
2 lysed as described above and the heme uptake was measure as described previously in 2.5.
3
4
5
6

7 **2.11. Statistical analysis.** Statistical analyses were conducted using GraphPad Prism 5
8 software (GraphPad Software, Inc., San Diego, CA). The data are presented as the mean \pm SD.
9 The experimental data were analyzed by one-way analysis of variance (ANOVA), and
10 differences between groups were assessed using Tukey's post-test. The level of significance
11 was set at $p < 0.05$, and all experiments were repeated at least three times.
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

3. Results

3.1 The effect of the heme analogues (SnPPIX, PdMPIX and ZnMPIX) on heme uptake and cell proliferation in *T. cruzi* epimastigotes.

The heme analogues used in this work have a similar chemical structure to heme. The mesoporphyrin derivatives have replaced the two vinylene groups of the original structure of protoporphyrin IX by two ethyl groups besides, they contain Pd(II) and Zn(II) as central ion (PdMPIX and ZnMPIX, respectively). The SnPPIX analogue have replaced only central metal ion but conserve the protoporphyrin ring structure. All three molecules are good fluorophores, and several have been used to study heme uptake by fluorescent microscopy and heme trafficking in different tissue types and organisms (Worthington *et al.*, 2001, Lara *et al.*, 2005; Rao *et al.*, 2005; and Sartonelo *et al.*, 2010), including our previous work where PdMPIX was used to follow by fluorescence microscopy heme internalization in *T. cruzi* epimastigotes (Lara *et al.*, 2007).

As a first approach to try to elucidate the process of heme transport in *T. cruzi* epimastigotes, we designed and optimized conditions to evaluate heme uptake in the presence of the fluorescent heme analogues SnPPIX, PdMPIX and ZnMPIX. The intracellular heme concentration was determined using hemochrome pyridine methods (Falk, 1964), as described in the materials and methods section. We observed that epimastigotes in the stationary phase can accumulate approximately 10 to 15 nmol of heme per 10^8 cells when they are incubated with 10 μ M heme for 10 minutes after being depleted of heme, as shown in Figure 1 (panels A, B and C; the concentration 0 represent cells without treatment with heme analogues). When the cells were treated with different concentrations of heme analogues, we observed a decrease in the intracellular heme concentration. This reduction is proportional to the analogue concentration, as shown in Figure 1 (A, B and C). In particular, we followed ZnMPIX uptake by spectrofluorometry, as described in the Materials and Methods. The fluorescence intensity

1 (IF) of ZnMPIX, which was measured from total cell extracts treated with different heme
2 concentrations, is presented in Figure 2. We observed lower ZnMPIX IF values as the heme
3 concentration increased. This reduction in ZnMP uptake is in agreement with the previous
4 concentration increased. This reduction in ZnMP uptake is in agreement with the previous
5 results presented in Figure 1, when we followed the intracellular heme uptake in parasites
6 treated with different concentrations of analogues. A similar effect was observed when
7 different *T. cruzi* strains were subject of the same treatment, such as CL-Brener and CL 14
8 (data not shown). These results indicate that heme analogues (PPIX or MPIX) might be
9 imported into cells through the same system used to uptake heme.
10

11 We also evaluated the effect of these heme analogues on the proliferation of *T. cruzi*
12 epimastigotes, as described in the material and methods. Briefly, we treated the epimastigotes
13 with heme and various concentrations of SnPPIX, PdMPIX or ZnMPIX. The parasites were
14 kept in culture for 12 days, and cell growth was monitored by counting in a Neubauer chamber.
15 We observed a marked inhibition in parasite growth when the parasites were treated with heme
16 analogues (Figure 3). However, the addition of heme after treatment with SnPPIX and PdMPIX
17 partially alleviated this inhibitory effect (Figures 3A and B). Interestingly, the inhibitory effect
18 observed after treatment with ZnMPIX did not revert in the presence of heme (Figure 3C),
19 suggesting that under these experimental conditions the effect of this compound might be more
20 deleterious or that ZnMPIX cannot be displaced by heme after being imported by the cells.
21 However, we cannot exclude other factors related with the experimental conditions and/or
22 related to any specific strain. We performed several controls to evaluated ZnMPIX toxicity in
23 CL-14 and Cl Brener epimastigotes and these assays did not show the deleterious effect of
24 ZnMPIX. In those cases the cells were treated during 10 min with (or without) 50 μ M ZnMPIX
25 in PBS buffer, then they were washed and cultured in BHI with 10% FCS (plus 30 μ M heme)
26 and we did not observe differences between the treated and non-treated cells (data not show).
27 More experiments are necessary to distinguish between these hypotheses about ZnMPIX
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 behavior compared to the others heme analogues. Besides, all the heme analogues showed a
2 deleterious effects when they replaced heme in the culture media. This result confirms that
3 heme is an essential cofactor for *T. cruzi* epimastigotes and these analogues may compete with
4 heme for the same carrier in the parasite cell.
5
6
7
8
9

10 **3.2 The effect of ABC transporter inhibitors on heme uptake and cell proliferation in *T.*** 11 ***cruzi* epimastigotes.** 12 13 14

15 Previously, we observed that the internalization of PdMPIX by epimastigotes, as measured by
16 fluorescence microscopy, decreased in the presence of ABC transporter inhibitors (Lara *et al.*,
17 2007). In addition, preliminary results obtained by J.A. Cricco and colleagues show that heme
18 uptake by epimastigotes is dependent on ATP (personal communication, data not shown).
19 These results suggest that heme uptake is mediated by an active transport system and suggest
20 that an ABC transporter is involved in this process. To elucidate what types of transporters or
21 carriers are involved in heme transport in epimastigotes, we evaluated the changes in
22 intracellular heme levels when parasites were treated with different transporter inhibitors. We
23 measured the intracellular heme level spectrophotometrically, as described before, to evaluate
24 the effect produced by the ABC transporter inhibitors cyclosporin A, indomethacin and
25 verapamil at concentrations of 10 μ M, 300 μ M and 2 μ M, respectively. The chosen inhibitors
26 and concentrations were based on our previous results (Lara *et al.*, 2007). We observed that
27 when parasites are treated with the inhibitors and then incubated with heme, less heme was
28 incorporated than in the untreated cells. The heme level decreased approximately 50% in all
29 cases, shown in Figure 4. This result supports our hypothesis that an ABC transporter could be
30 involved in the translocation of heme through the lipid bilayer in *T. cruzi*. To evaluate the effect
31 of these inhibitors on the parasite proliferation, we treated them with increasing doses of
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

cyclosporin A (1 μM to 20 μM). It was observed that doses smaller than 5 μM do not interfere with proliferation. At concentration of 10 μM CsA we observed a small reduction on parasite proliferation. However, at cyclosporin A concentrations greater than 10 μM , the proliferation induced by heme incubation is prevented, as shown in Figure 5. Finally, we decided to evaluate how increasing doses of heme (10 μM to 300 μM) modulate the effect of 10 μM cyclosporin A treatment. As shown in Figure 6, our results demonstrate that heme does not reverse the inhibitory effects of cyclosporin A, suggesting that CsA, at this concentration, is affecting other processes in addition to heme uptake. To further investigate this point, we evaluated the intracellular heme level in the presence of CsA.

Parasites were treated with 10 μM cyclosporin A for 30 minutes and then incubated with or without heme in culture for 7 days, as detailed in the material and methods. These results are shown in Figure 7. We observed that CsA treatment prevents heme uptake when the parasites are incubated with 300 μM heme. However, a basal level of heme was maintained in the cells, this amount was presumably sufficient to allow the parasite regular proliferation. In addition, the results presented in Figure 6 and 7 allowed us consider that at 10 μM CsA the parasite proliferation might be affected because extra heme import is prevented, inhibiting at least in part, the heme uptake processes

4. Discussion

The study of heme acquisition by *T. cruzi* is critical for understanding the following aspects of *T. cruzi* infection: i) the biology of the parasite, ii) heme uptake in *T. cruzi*, and iii) the interaction between the parasite and the vector. Advancing knowledge regarding the transporters used by parasites is necessary because blocking heme transport may interrupt the relationship between *T. cruzi* and the vector or host. In this work, we developed a method to evaluate the heme uptake processes in *T. cruzi* epimastigotes. We used the fluorescent heme

1 analogues SnPPIX, ZnMPIX and PdMPIX to evaluate heme uptake. We observed that the
2 presence of these compounds decreased intracellular heme levels and affected parasite
3 proliferation, suggesting a competition between the porphyrins and heme for the same transport
4 machinery. The presence of common porphyrin-transporter in *T. cruzi*, is in agree with the
5 results presented by Campos-Salinas *et al.* were they study the heme transport by *Leishmania*
6 *donovani* promastigotes using MgPPIX as heme analogue by flow citometry (Campos-Salinas,
7 2011). Our results indicate that *T. cruzi* proliferation is mediated almost exclusively by heme
8 through unknown mechanisms. We observed that treatment with SnPPIX, ZnMPIX and
9 PdMPIX inhibited parasite proliferation, and this inhibition is reverted by the presence of
10 heme, although the case of ZnMPIX is controversial and more studies will be necessary to
11 address it behavior.

12 ABC transporters are responsible for moving a variety of molecules through the lipid
13 membrane of cells and organelles, mostly against a concentration gradient at the expense of
14 ATP hydrolysis. These carriers are remarkably variable with regards to their physiological
15 functions, their substrates and the direction of transport, alternating between import and export
16 (Higgins, 1992). Klokouzas and colleagues identified approximately 30 members of the ABC
17 family in the main protozoan parasites (Klokouzas *et al.*, 2003). Sauvage and colleagues
18 presented a review on ABC proteins that have been identified in the main protozoan parasites,
19 including *Leishmania*, *Trypanosoma*, *Plasmodium*, *Toxoplasma*, *Cryptosporidium* and
20 *Entamoeba* (Sauvage *et al.*, 2009). It has also been observed that the activity of ABC
21 transporters is a basic biological strategy to defend living cells against cytotoxic attack by
22 xenobiotics, but ABC transporters also play an important role in other cellular functions that
23 have yet to be characterized in protozoan parasites. In the present research work, we studied the
24 effects of specific ABC transporter inhibitors (cyclosporin A, verapamil and indomethacin) to
25 investigate the possibility for the involvement of a protein of this family in the translocation of

1 heme across the plasma membrane of *T. cruzi* epimastigotes. We showed that heme uptake by
2 the parasite decrease approximately 50% in the presence of these inhibitors. In subsequent
3 analyses, we evaluated the effect of cyclosporin A on parasite proliferation and heme
4 accumulation. It was possible to observe a reduction in cell proliferation concomitantly to a
5 reduction on heme accumulation. These results allow us to hypothesize that interference with
6 heme transport will interrupt the trypanosome life cycle. The inhibition of *Trypanosoma cruzi*
7 cell proliferation by cyclosporin A was also observed by Búa and colleagues (Búa *et al.*, 2008).
8 The authors reported inhibitory effects of cyclosporin A on the proliferation of epimastigotes,
9 the internalization of trypomastigotes, the development of intracellular amastigotes, and
10 infection by *T. cruzi* *in vivo*. They associate the inhibitory effect of CsA with a role of a Pgp
11 protein, but this was unrelated to the transport of heme and/or porphyrins. The involvement of
12 ABC transporters in heme transport has been observed in other studies. Krishnamurthy and
13 colleagues published a study demonstrating the involvement of an ABC transporter in the
14 translocation of coproporphyrinogen III (CPIII), a precursor of heme, from the cytoplasm into
15 the mitochondria in mammalian cells by linking the expression of this transporter with heme
16 synthesis (Krishnamurthy *et al.*, 2006). Other studies in mammals have shown that ABCG2
17 (Krishnamurthy *et al.*, 2004) has a potential heme export activity during the development of
18 erythroid cells. Yet another example can be seen in the recent study by Desuzinges-Mandon
19 and colleagues (Desuzinges-Mandon *et al.*, 2010), which demonstrates the ability of the
20 membrane protein ABCG2 to transfer the heme molecule to human serum albumin. Recently,
21 an ABC protein was shown to be involved in internal heme trafficking in *Leishmania donovani*
22 (Campos-Salinas *et al.*, 2011). This protein is utilized for hemoglobin-bound heme trafficking
23 to the mitochondria and not in the movement of free heme. This protein is conserved in other
24 trypanosomatids. Other studies presented by Vanhollebeke and colleagues demonstrated that
25 TbHpHbR is a bloodstream stage-specific protein and directs the internalization of heme
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 carried by the Hp-Hb complex into hemoproteins in order to optimize growth of bloodstream
2 forms (Vanhollebeke *et al.*, 2008). TbHpHbR is present in *T. brucei subsp. gambiense* and *T.*
3
4 *brucei subsp. rhodesiense* but absent from the related kinetoplastids *T. cruzi* and *Leishmania*. It
5
6 is important to note that although only 3 ABC genes have been characterized in *T. cruzi*, 28
7
8 have been identified in the genome of this parasite; the functions of the remaining 25 are still
9
10 unknown (Sauvage, 2009).
11
12

13
14 In summary, our results indicate that a common porphyrin transporter may be involved
15
16 in the entry of amphipathic porphyrins, as suggested by the observation that different heme
17
18 analogues compete with heme for cellular uptake. We present evidence indicating that
19
20 heme/porphyrins translocation in *T. cruzi* epimastigote is mediated by a transporter protein and
21
22 possibly a member of the ABC transporter family; however, a molecular approach for
23
24 identifying the transporter is required. The answer to this question in *T. cruzi* will be of great
25
26 importance because heme transport is essential for parasite survival. The identification of this
27
28 essential transporter might allow for the development of chemotherapy targets for the treatment
29
30 of Chagas' disease.
31
32
33
34
35
36
37
38

39 **5. Acknowledgement**

40 This work was supported by Coordenação de Aperfeiçoamento de Pessoal de Nível
41
42 Superior (CAPES), Fundação de Amparo a Pesquisa do Estado do Rio de Janeiro (FAPERJ) and
43
44 Instituto Nacional de Ciência e Tecnologia - Entomologia Molecular (INCT-EM) – Brasil to
45
46 MCP. PIP 2010-0685 Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET)
47
48 Argentina and Subsidio de la Fundación Bunge y Born 2010/2012, para Investigación de la
49
50 Enfermedad de Chagas (FBB27/10) to JAC.
51
52
53

54 **6. References**

55
56
57
58 Búa, J., Fichera, L.E., Fuchs, A.G., Potenza, M., Dubin, M., Wenger, R.O., Moretti, G.,
59
60 Scabone, C.M., Ruiz, A.M., 2008. Anti-*Trypanosoma cruzi* effects of cyclosporin A
61
62
63
64
65

1 derivatives: possible role of a P-glycoprotein and parasite cyclophilins. *Parasitology*. 135, 217-
2 228.

3
4
5 Campos-Salinas, J., Cabello-Donayre, M., García-Hernández, R., Pérez-Victoria, I., Castanys,
6 S., Gamarro, F., Pérez-Victoria, J. M., 2011. A new ATP-binding cassette protein is involved in
7 intracellular haem trafficking in *Leishmania*. *Mol. Microbiol.*79, 1430-1444.

8
9
10
11
12 Coura J.R., Viñas P.A., 2010. Chagas disease: a new worldwide challenge. *Nature*. 465, S6-S7.

13
14
15
16 Desuzinges-Mandon, E., Arnaud, O., Martinez, L., Huche, F., Di Pietro, A., Falson, P., 2010.
17 Abcg2 transports and transfers heme to albumin through its large extracellular loop. *J. Biol.*
18 *Chem.* 285, 33123-33.

19
20
21
22
23 Deterding, L.J., Ramirez, D.C., Dubin, J.R., Mason, R.P., Tomer, K.B., 2004. Identification of
24 free radicals on hemoglobin from its self-peroxidation using mass spectrometry and immuno-
25 spin trapping: observation of a histidinyl radical. *J. Biol. Chem.* 279, 11600-11607.

26
27
28
29
30
31 El-Sayed, N.M., Myler, P.J., Bartholomeu, D.C., Nilsson, D., Aggarwal, G., Tran, A.N., et al.,
32 2005. The genome sequence of *Trypanosoma cruzi*, etiologic agent of Chagas disease. *Science*.
33 309, 409-15.

34
35
36
37
38 Falk, J.E., 1964. *Porphyryns and Metalloporphyryns*. Amsterdam, Elsevier Publishing
39 Company.

40
41
42
43
44 Graça-Souza AV, Maya-Monteiro C, Paiva-Silva GO, Braz GRC, Paes M C et al., 2006.
45
46 Adaptations against heme toxicity in blood-feeding arthropods. *Insec Bioch Mol Biol* 36: 322-
47
48 335.

49
50
51
52
53 Higgins, C.F., 1992. ABC transporters: from microorganisms to man. *Annu. Rev. Cell. Biol.* 8,
54 67-113.

1 Hotez, P.J., Molyneux, D.H., Fenwick, A., Kumaresan, J., Sachs, S.E., Sachs, J.D., Saviolli, L.,
2 2007. Control of Neglected Tropical Diseases. *N. Engl. J. Med.* 357, 1018-27.

3
4
5 Klokouzas, A., Shahi, S., Hladky, S.B., Barrand, M.A., Van Veen, H.W., 2003. ABC
6 transporters and drug resistance in parasitic protozoa. *Int. J. Antimicrob. Agents.* 22, 301–17.

7
8
9
10 Koreny, L., Lukes, J., Oborník, M., 2010. Evolution of the haem synthetic pathway in
11 kinetoplastid flagellates: An essential pathway that is not essential after all? *Int. J. Parasitol.* 40,
12 149-56.

13
14
15
16
17
18 Krishnamurthy, P. C., Du, G., Fukuda, Y., Sun, D., Sampath, J., Mercer, K. E., Wang, J., Sosa-
19 Pineda, B., Murti, K. G., Schuetz, J. D., 2006. Identification of a mammalian mitochondrial
20 porphyrin transporter. *Nature.* 443, 586–589.

21
22
23
24
25 Krishnamurthy, P., Ross, D. D., Nakanishi, T., Bailey-Dell, K., Zhou, S., Mercer, K. E.,
26 Sarkadi, B., Sorrentino, B. P., Schuetz, J. D., 2004. The stem cell marker Bcrp/ABCG2
27 enhances hypoxic cell survival through interactions with heme. *J. Biol. Chem.* 279, 24218–
28 24225.

29
30
31
32
33
34 Lara, F.A., Lins, U., Bechara, G.H., Olivera, P.L., 2005. Tracing heme in a living cell:
35 hemoglobin degradation and heme traffic in digest cells of the cattle tick *Boophilus microplus*.
36 *J. Exp. Biol.* 208, 3093-3101.

37
38
39
40
41
42 Lara, F.A., Sant'Anna, C., Lemos, D., Laranja, G.A.T., Coelho, M.G.P., Reis Salles, I., Michel,
43 A., Oliveira, P.L., Cunha-e-Silva, N., Salmon, D., Paes, M.C., 2007. Heme requirement and
44 intracellular trafficking in *Trypanosoma cruzi* epimastigotes. *Biochem. Biophys. Res.*
45 *Commun.* 355, 16-22.

46
47
48
49
50
51 Lombardo, M.E., Araújo, L.S., Batlle, A., 2003. 5-Aminolevulinic acid synthesis in
52 epimastigotes of *Trypanosoma cruzi*. *Int. J. Biochem. Cell Biol.* 35, 1263-1271.

53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

1 Rao, A.U., Carta, L.K., Lesuisse, E., Hamza, I., 2005. Lack of heme synthesis in a free-living
2 eukaryote. Proc. Natl. Acad. Sci. 102, 4270- 4275.

3
4
5 Ryter, S.W., Tyrrel, R.M., 2000. The heme synthesis and degradation pathways: role in oxidant
6 sensitivity: haem oxygenase has both pro and antioxidant properties. Free Radic. Biol. Med. 28,
7 289-309.
8
9

10
11
12 Sartonello, R., Budu, A., Bagnaresi, P., Fernandes, C.A.H., Sato, P.M., Bueno, V.B., Fontes,
13 M.R.M., Oliveira, P.L., Paiva-Silva, G.O., Alves, S.V., Netto, L.E.S., Catalani, L.H., Garcia,
14 C.R.S., 2010. *In vivo* uptake of a heme-analogue Zn-Protoporphyrin IX by the human malaria
15 parasite *P. falciparum*-infected red blood cells. Cell Biol. Int. 34, 859-65.
16
17
18
19

20
21
22 Sauvage, V., Aubert D., Escotte-Binet, S., Villena, I., 2009. The role of ATP-binding cassette
23 (ABC) proteins in protozoan parasites. Mol. Biochem. Parasitol. 167, 81–94.
24
25

26
27 Severance, S., Hamza, I., 2009. Trafficking of Heme and Porphyrins in Metazoa. Chem. Rev.
28 109, 4596–4616.
29
30

31
32 Tripodi, K.E.J., Menendez Bravo, S.M., Cricco, J.A., 2011. Role of heme and heme-proteins
33 trypanosomatid essential metabolic pathways. Enzyme Research 2011, 873230. Epub 2011 Apr
34 10.
35
36
37

38
39
40 Vanhollebeke, B., De Muylder, G., Nielsen, M.J., Pays, A., Tebabi, P., Dieu, M., Raes, M.,
41 Moestrup, S.K., Pays, E., 2008. A haptoglobin-hemoglobin receptor conveys innate immunity
42 to *Trypanosoma brucei* in humans. Science 320, 677-681.
43
44
45

46
47 Wigglesworth VB., 1943. The fate of haemoglobin in *Rhodnius prolixus* (Hemiptera) and other
48 blood-sucking arthropods. Proc.R.Soc. London 131, 313–339
49
50

51
52
53 World Health Organization. Fact sheet n° 340. June 2010 from
54 <http://www.who.int/mediacentre/factsheets/fs340/en/print.html>.
55
56
57
58
59
60
61
62

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Worthington M.T., Cohn S.M., Miller S.K., Luo R.Q., Berg C.L., 2001. Characterization of a human plasma membrane heme transporter in intestinal and hepatocyte cell lines. *Am J Physiol Gastrointest Liver Physiol* 280, G1172-1177.

Accepted Manuscript

7. Legends

1
2
3
4 **Figure 1. The heme analogues (SnPPIX, PdMPIX and ZnMPIX) prevents heme uptake**
5
6 **by *T. cruzi* epimastigotes.** *Trypanosoma cruzi* Dm28c were maintained in BHI supplemented
7
8 with 10% FBS at 28°C for 7 days. The epimastigotes were washed with PBS and treated with
9
10 different concentrations of SnPPIX (A), PdMPIX (B) and ZnMPIX (C) for 10 min and then
11
12 incubated with 10 µM heme for 10 min. The negative control (0) did not receive any treatment,
13
14 whereas the positive control (black bar) was treated with 10 µM heme for 10 min. Heme uptake
15
16 in *T. cruzi* was measured in lysed epimastigotes using the alkaline pyridine method. All data
17
18 are presented as the means ± SD, *p < 0.05 in relation to the heme treatment, as determined
19
20 using Tukey's test. This result is representative of 3 experiments.
21
22
23
24
25
26
27

28
29 **Figure 2. The presence of increased heme concentration affects the ZnMPIX uptake in *T.***
30 ***cruzi* epimastigotes.** *T. cruzi* Dm28c epimastigotes were maintained in BHI supplemented with
31
32 10% FCS and 30 µM Heme at 28°C for 7 days. The cells were washed twice with PBS and
33
34 suspended in the same buffer at 2×10^8 cells/ml. The cells were incubated with ZnMPIX and
35
36 different concentrations of heme for 5 min before the reaction was stopped by the addition of a
37
38 cold PBS solution containing 1 mM heme. The amount of uptake was measured in lysed
39
40 epimastigotes, after several steps of centrifugation and pellet suspension in cold PBS buffer, by
41
42 determining fluorescence intensity in a spectrofluorometer ($\lambda_{\text{ex}} = 405$ nm and $\lambda_{\text{EMmax}} = 577$
43
44 nm). All data are presented as the means ± SD. This result is representative of 3 experiments.
45
46
47
48
49
50
51
52
53

54 **Figure 3. The heme analogues (SnPPIX, PdMPIX and ZnMPIX) negatively affect the**
55 **growth kinetics of *T. cruzi* epimastigotes.** *Trypanosoma cruzi* of the Dm28c strain were
56
57 maintained at 28°C in BHI supplemented with 10% FBS and 30 µM heme. Epimastigotes in the
58
59
60
61
62
63
64
65

1 stationary phase were washed with PBS and incubated in BHI supplemented with 10% FBS.
2 The epimastigotes were then treated with different concentrations of SnPPIX (A), PdMPIX (B)
3 and ZnMPIX (C) for 10 min before incubation with 30 μ M heme. All data are presented as the
4 means \pm SD, *p < 0.05 in relation to the heme treatment by Tukey's test. The results are
5 representative of three independent experiments.
6
7
8
9
10

11 **Figure 4. The treatment with ABC transporter inhibitor affects the heme uptake in *T.***
12 ***cruzi* epimastigotes.** *Trypanosoma cruzi* Dm28c were maintained in BHI supplemented with
13 10% FBS at 28°C for 7 days. The epimastigotes were washed with PBS and treated with 10 μ M
14 CsA, 300 μ M indomethacin or 2 μ M verapamil for 30 min and then incubated with 10 μ M
15 heme for an additional 10 min. The control did not receive any treatment. Heme uptake in *T.*
16 *cruzi* was measured in lysed epimastigotes using the alkaline pyridine method. All data are
17 presented as the means \pm SD of three independent experiments, *p < 0.05 in relation to the
18 heme treatment, as measured by Tukey's test.
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35

36 **Figure 5. The dose-effect of CsA on the proliferation of *T. cruzi* epimastigotes grown in**
37 **the presence of heme.** *Trypanosoma cruzi* Dm28c were maintained at 28°C in BHI
38 supplemented with 10% FBS and 30 μ M heme. Epimastigotes in the stationary phase were
39 washed with PBS and treated with different concentrations of cyclosporin A (CsA) for 30 min.
40 The parasites were then incubated with 10 μ M heme for 7 days. All data are presented as the
41 means \pm SD of two independent experiments, *p < 0.05 in relation to the heme treatment, as
42 determined using Tukey's test.
43
44
45
46
47
48
49
50
51
52
53
54


55 **Figure 6. The dose-effect of heme on the proliferation of *T. cruzi* epimastigotes grown in**
56 **the presence of CsA.** *Trypanosoma cruzi* Dm28c were maintained in BHI supplemented with
57 10% FBS and 30 μ M heme at 28°C. Epimastigotes in the stationary phase were washed with
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

PBS and treated with 10 μM cyclosporin A (CsA) for 30 min. The cells were then incubated with different concentrations of heme for 7 days. All data are presented as the means \pm SD of two independent experiments. * $p < 0.05$ in relation to the cells grown with 30 μM heme, and $^+p < 0.05$ in relation to the cells grown with 300 μM heme, as determined using Tukey's test.

Figure 7. The effect of cyclosporin A on heme uptake in *T. cruzi* epimastigotes.

Trypanosoma cruzi Dm28c were maintained at 28°C in BHI supplemented with 10% FBS and 30 μM heme. Epimastigotes in the stationary phase were washed with PBS and treated with 10 μM cyclosporin A (CsA) for 30 min. The cells were then incubated with 300 μM heme for 7 days. Heme uptake in *T. cruzi* was measured in lysed epimastigotes using the alkaline pyridine method. All data are presented as the means \pm SD with * $p < 0.05$ in relation to the heme treatment, as determined using Tukey's test. This result is representative of two independent experiments.



Accepted Manuscript






Fig. 2. Cupello MP *et al.*, 2011


A



B



C

Fig. 3. Cupello MP *et al.*, 2011




Fig. 4 Cupello MP *et al.*, 2011




Fig. 5. Cupello MP *et al.*, 2011




Fig. 6. Cupello MP *et al.*, 2011




Fig. 7. Cupello MP *et al.*, 2011