

Effect of cholesterol reduced and zinc fortification treatments on physicochemical, functional, textural, microstructural and sensory properties of soft cheese

MICAELA GALANTE^a, YANINA PAVÓN^b, SANDRA LAZZARONI^b, MARINA SOAZO^{a,c}, SILVIA COSTA^b; VALERIA BOERIS^{a,d}, PATRICIA RISSO^{a,e,f*}, SERGIO ROZYCKI^b

^aFacultad de Ciencias Bioquímicas y Farmacéuticas-CONICET, Universidad Nacional de Rosario (UNR), Suipacha 531, 2000 Rosario, Argentina.

^bInstituto de Tecnología de Alimentos, Facultad de Ingeniería Química, Universidad Nacional del Litoral (UNL), Santiago del Estero 2829, 3000 Santa Fe, Argentina.

^cInstituto de Química Rosario-CONICET, Suipacha 531, 2000 Rosario, Argentina.

^dFacultad de Química e Ingeniería, Pontificia Universidad Católica Argentina, Pellegrini 3314, 2000 Rosario, Argentina.

^eInstituto de Física Rosario-CONICET, 27 de Febrero 210bis, 2000 Rosario, Argentina.

^fFacultad de Ciencias Veterinarias, UNR, Ovidio Lagos y Ruta 33, 2170 Casilda, Argentina.

Running headline: Evaluation of functional soft cheese

Corresponding author: Risso, Patricia

Phone number: +54 (0341) 4804597 int. 253

e-mail address: phrisso@yahoo.com.ar

Postal address: Suipacha 531, 2000 Rosario, Argentina

Abstract

The aim of this study was to produce and evaluate a soft cheese fortified with zinc and with cholesterol-reduced content. To meet this objective a cream base was prepared, from which Cholesterol was removed using β -cyclodextrin as extracting agent. Then, cholesterol-reduced content cheese with and without the addition of zinc salts (ZnSO_4 or ZnCl_2) was produced. Additionally, a cheese without any treatment was prepared. Furthermore, physicochemical, textural, functional, microstructural and sensory determinations were performed. As a result, 87-94% zinc fortified and 93% cholesterol-reduced cheese samples were obtained which had similar sensorial characteristics to the cheese without treatment.

Keywords: Soft cheese, Zinc Fortification, Cholesterol removal, β -cyclodextrin

INTRODUCTION

In the last years, the “functional foods” concept has achieved a great importance for consumers and between them dairy products are the most required. It has been reported that high cholesterol blood levels represent a major risk factor for developing coronary heart disease. In spite of the relationship between dietary cholesterol and plasma cholesterol levels it is being still investigated, the World Health Organization and the American Heart Association have recommended the reduction of dietary cholesterol consumption (Dias *et al.* 2010). Reduce cholesterol from fat dairy products is a way to avoid the risk of these diseases with keeping the rest of the milk fat, since is an excellent source of energy, vitamins (A, D and E), antioxidants and essential fatty acids (Rehm *et al.* 2015). In addition, it is important to highlight that fat removal from dairy products affects in a negative way their sensorial characteristics (Kim *et al.* 2009).

The methods to reduce cholesterol from dairy products include physical, chemical or biological strategies, between them could be mentioned the extraction with high-methoxyl pectins, plant sterols, adsorption with saponin and solvent. However, most of these methods are not suitable, since other molecules are removed at the same time than the cholesterol (Dias *et al.* 2010). The use of β -cyclodextrin (β -CD) is a strategy that effectively removes cholesterol from dairy products improving their nutritional characteristics, leaving the rest of milk fat. This method is selective and avoids the removal of flavours and nutritional components during the cholesterol extraction. β -CD has a central cavity which allows complex formation with the cholesterol molecule (Kwak *et al.* 2001, Kwak *et al.* 2002). In addition, β -CD is nontoxic, edible, non-hygroscopic, chemically stable and easy to separate. Furthermore, in 1998 it was introduced into the Generally Recognized As Safe list of compounds (GRAS), because it is not absorbed in the upper gastrointestinal tract, and it is completely metabolized by the colon microflora (Dias *et al.* 2010). Moreover, recent works have shown that the use of β -CD as a strategy to remove cholesterol from dairy products does not show significant effects on their chemical, rheological, and sensory properties (Dias *et al.* 2010). All these positive attributes make β -CD use a suitable strategy for cholesterol removal from food.

Zinc is one of the most important trace elements present in the body with a great nutritional importance and that has a recognized action on more than 300 enzymes, participating in either their structure or in their catalytic and regulatory action. Due to the great number of zinc-dependent biological processes, the deficiency states of this essential nutrient entails multiclinical consequences, which range from mild to serious dysfunctions (Salgueiro *et al.* 2000, Pomastowski *et al.* 2014).

Zn^{2+} deficiency states emerge as a consequence of two groups of causes. The first group includes several syndromes related to metabolic or genetic malfunctions and the second one comprises nutritional causes like decreased Zn^{2+} intake or consumption of poor Zn^{2+} content foods, which represent the most important and the most common (Salgueiro *et al.* 2000). The Recommended Dietary Allowance (RDA), according to the Food and Agriculture Organization of the United Nations and the World Health Organization (FAO/WHO), for Zn^{2+} is 3.9 mg day^{-1} for females and 4.2 mg day^{-1} for males, assuming a bioavailability of dietary zinc of 50 % (diet high in meat), and 9.8 mg day^{-1} for females and 14 mg day^{-1} for males, assuming a bioavailability of dietary zinc of 15 % (vegetarian diets) (WHO/FAO 2005, Kahraman and Ustunol 2012).

The available studies clearly show that food fortification with Zn^{2+} can increase the dietary intake and the total daily absorption of this trace element (Hess and Brown 2009). In addition, is extremely important to control the quantity of mineral supplementation in food to ensure the desired level in the final products does not exceed the critical limits for an adult man, which is 45 mg/day according to FAO/WHO (WHO/FAO 2005). The selection of an adequate vehicle is crucial for successful fortification. Milk and dairy products are a good option for Zn^{2+} fortification, not only due to their worldwide consumption, but also because of their high nutritional value, the buffer effect in the digestion and absorption processes and the low pH value (in fermented products) that enhances the Zn^{2+} solubilization increasing its bioavailability (Aquilanti *et al.* 2012). Zinc sulfate and zinc chloride salts belong to the list of compounds listed as GRAS (Kahraman and Ustunol 2012).

Previous works present strategies for Zn^{2+} fortification of different kinds of cheese and dairy products, thus supporting the use of fortification with Zn^{2+} as a method to counteract intake deficiency (El-Din *et al.* 2012, Kahraman and Ustunol 2012).

Cuartirollo cheese is a soft cheese defined by the Código Alimentario Argentino as a high (46-55%) or very high (>55%) moisture, fatty (45-60%), manufactured with raw milk, acidified by lactic cultures and coagulated by rennet or specific enzymes (ANMAT 2002). The treatments (homogenisation, heating) and the addition of different cosolutes (β -CD, zinc salts) in order to cholesterol-removal and Zn^{2+} fortification may alter dairy products characteristics. The aim of this work is to evaluate physicochemical, sensory, functional, rheological, textural and microstructural characteristics of a cholesterol-reduced and Zn^{2+} fortified Cuartirolo type soft cheese.

MATERIALS AND METHODS

Materials

The Cream (48% fat content) and the skim milk (< 0.05 % w/w fat content) used to obtain the cream base (CB) were purchased from Manfrey (Santa Fe, Argentina). Commercial β -CD (Kleptose®, Roquette, France) was the extracting agent used to remove cholesterol from the CB. The fortification agents, $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ and ZnCl_2 (anhydrous), were purchased from Novalquim (Santa Fe, Argentina) and Cicarelli (Santa Fe, Argentina), respectively. Salts belong to the list of compounds listed as GRAS.

Methods

Cholesterol-reduced and Zn^{2+} fortified Cuartirolo soft cheese manufacture

Preparation of cholesterol-reduced content cream base

The quantities of cream (48 % w/w fat content) and skim milk (< 0.05 % w/w fat content) required to obtain a cream base (CB: 10 % w/w fat content) were mixed and homogenised using a two stages homogeniser ST2 (Simes S.A., Santa Fe, Argentina) with a total pressure of 100 atm at 70 °C in order to promote cholesterol extraction.

The homogenised CB was brought to 30 °C, where β -CD was added at a ratio of 2.85 % w/w after which the final mixture was left in mechanic agitation for 30 min. The system was centrifuged at 700 g (Mistral 4L, MSE, Crawley, UK) and 20 °C for 15 min in order to precipitate the β -CD/cholesterol complex, obtaining a CB with cholesterol-reduced content in the supernatant. The optimal conditions for cholesterol extraction were determined using a response surface methodology, which was performed in an early stage (data not shown).

Cheese-making process

The cheese samples were made according to traditional Cuartirolo technology. A bench-top (< 10 L) cheese vat was used. The cheese-making process was made in duplicated for each type of cheese in study.

Each type of cheese was made from CB treated with β -CD and diluted with skim milk (Milkaut S.A., Santa Fe, Argentina) to lower down the fat content to 3 % w/w (treated milk, TM), which was pasteurized at 65 °C for 20 min. This TM was then cooled to 40 \pm 1 °C, followed by the addition of CaCl₂ (200 mg Kg⁻¹, Cicarelli, Santa Fe, Argentina) and the Zn²⁺ salts (16 mg Zn²⁺ kg⁻¹) in the case of fortified cheese samples (Aquilanti *et al.* 2012, Kahraman and Ustunol 2012). The starter lactic culture for Cuartirolo soft cheese (STD, Diagramma S.A., Santa Fe, Argentina) was added at a ratio of 0.01 g kg⁻¹ of TM. After that, bacterial rennet (0.94 UR mL⁻¹, Chy-Max®, Chr.Hansen, Denmark) was added at a ratio of 0.5 mL kg⁻¹ of TM. Clot was formed after 30 min, cubes of approximately 2 cm³ were cut and allowed to heal during 15 min.

The mixture whey-curd particles were gently stirred for 5 min. This operation repeated thrice. Whey was drained and the curd was moulded. Moulded cheese (0.5 kg cheese) were stored at 40 \pm 1 °C, inverted each every 30 min until the pH of the cheese was near 5.3 \pm 0.1. The molded cheese was maintained in a cold brine solution (20 %

w/w NaCl) for a period of 1 h kg⁻¹ of cheese. After that, cheese samples were vacuum packed and ripened at 5 °C for 20 days (Figure 1).

Four types of Cuartirolo soft cheese were made: 1) a cholesterol-reduced cheese (ChC-R: made of TM obtained from cholesterol-reduced CB and without Zn²⁺ fortification), 2) a ZnCl₂ fortified cheese (ChZnCl₂: made of TM obtained from cholesterol-reduced CB and fortified with ZnCl₂ salt), 3) ZnSO₄ fortified cheese (ChZnSO₄: made of TM obtained from cholesterol-reduced CB and fortified with ZnSO₄ salt), and 4) a cheese without any treatment (ChWT: made without cholesterol extraction treatment and without fortification). Each cheese sample was made in duplicated.

Quantification of cholesterol

Cholesterol content of the CB, before and after β-CD treatment, was determined in quadruplicate. Samples of CB were saponified, followed by cholesterol extraction using n-hexane (Cicarelli, Santa Fe, Argentina) according to the technique published by Pavón *et al.* (2014). Cholesterol quantification was made by an enzymatic method using a commercial kit (Wiener Lab., Argentina). Cholesterol content of all samples was determined by absorption at 510 nm using a UV-VIS spectrophotometer (Jasco V550, Japan) and was compared with the cholesterol standard solution (belong to the commercial kit). Cholesterol extraction percentage (%ExtChol) was calculated as follows:

$$\%ExtChol = 100 - [(amount\ of\ cholesterol\ in\ \beta\text{-}CD\ treated\ CB * 100) / (amount\ of\ cholesterol\ in\ untreated\ CB)] \quad (1)$$

Chemical cheese analysis and cheese yield determination

Zinc content determination

Cheese Zn^{2+} content was determined according to the flame atomic absorption spectrophotometric method of the Association of Official Analytical Chemists (1990). The values corresponding to the percentage of Zn^{2+} fortification (% Zn^{2+} Fortification) are calculated as follows:

$$\% \text{Zn}^{2+} \text{ Fortification} = \frac{([\text{Zn}^{2+}]_{\text{ChF}} - [\text{Zn}^{2+}]_{\text{ChC-R}})}{[\text{Zn}^{2+}]_{\text{max}}} * 100 \quad (2)$$

where $[\text{Zn}^{2+}]_{\text{ChF}}$ and $[\text{Zn}^{2+}]_{\text{ChC-R}}$ are the Zn^{2+} concentration in the fortified cheese and the ChC-R respectively. $[\text{Zn}^{2+}]_{\text{max}}$ is the Zn^{2+} maximum concentration present in the cheese if all the Zn^{2+} salt added remains in the curd.

Cheese yield

Cheese yield was determined as the weight of curd obtained per 100 g of TM used for cheese-making, and the yield percentage (% Yield) according to:

$$\% \text{Yield} = \frac{(\text{wt.cheese} * 100)}{(\text{wt.milk})} \quad (3)$$

where wt.cheese and wt.milk are the weight of cheese and milk respectively (Kwak *et al.* 2001, Kwak *et al.* 2002).

Moisture content

For cheese moisture content determinations, 2 g of shredded cheese samples were weighted onto porcelain dishes and dried at 105 ± 1 °C until constant weight was reached, according to the Association of Official Analytical Chemists official method (1990).

Measurement of cheese firmness

Cylindrical samples of cheese (height: 20 mm, diameter: 40 mm) were prepared and tempered at room temperature for 1 h before the determinations. The firmness of the cheese was measured using a motorized test frame Mecmesin Multitest 2.5-d

(Mecmesin, Spain) equipped with a dynamometer of a 100 N load cell. Penetration tests were carried out in the middle of the cylindrical samples, always at the same place, to avoid side effects.

The force in Newton required to push a cylinder probe (diameter 20 mm), at a speed of 10 mm min⁻¹, into the cheese mesh was measured. Cheese firmness was considered as the force required to insert the probe to a depth of 10 mm into the cheese.

Colorimetric measurement

A high-resolution digital camera (Nikon, Coolpix P520) was used to measure colour by capturing the colour image of cheese samples under proper lighting. The digital images were processed, using Photoshop software (Adobe Systems Inc., USA) according to the method proposed by Soazo *et al.* (2014) in order to obtain the L*, a* and b* parameters. Where L* is the luminance or lightness component that goes from 0 (black) to 100 (white), and parameters a* (green to red) and b*(blue to yellow) are both chromatic components, varying from -120 to +120.

Microstructure analysis

Confocal scanning laser microscopy (CSLM)

CSLM is widely used for the study of food microstructure. Sections of samples, approximately of 5 mm X 5 mm X 2 mm, were cut with a scalpel placed in a microscope slide and one drop (50 µL) of 0.01 mg mL⁻¹ Rhodamine B (Sigma-Aldrich, St. Louis, USA) was added to the cut surface. Rhodamine B binds to proteins fraction in the cheese samples, labelling them in red.

Images of representative areas of each sample were captured using a confocal microscopy (Nikon Eclipse TE-2000-E, Japan), with an objective magnification of 60X (oil immersion lens) and a numerical aperture of 1.4.

The digital images were acquired with a pixel resolution of 1024 X 1024 and stored in a TIFF format in order to be analyzed.

Images textural analysis

Texture parameters: Shannon entropy (S), smoothness (K), uniformity (U) and grey level variance ($\sigma^2(N)$), were determined from the CLSM images according to the method proposed by Ingrassia *et al.* (2013).

Pore diameter determination

In order to determine cheese pore diameter from the CSLM images a plug-in of the image J (version 1.485) program was employed (Doube *et al.* 2010). From the CSLM images, the thickness parameter was obtained using the Bone J plug-in (version 1.3.12). This can be interpreted as the width of cavities or as the distance between structures. It can thus be used to estimate parameters such as pore diameter.

Cheese meltability determinations

A modified Schreiber's test was used to determine the fusion capability of cylindrical samples of cheese (diameter: 37 mm, height: 12 mm) (Muthukumarappan *et al.* 1999).

All cheese meltability determinations were made after 20 days of ripening. Samples were maintained at 4 °C for 30 min before the determinations and then were placed in a stove at 130 °C for 15 min. Subsequently, both areas (before and after heating) were determined in order to quantify the fusion capability of each cheese sample.

Sensory analysis

Cheese samples were evaluated by a trained sensory panel composed of ten members (6 female and 4 male from 25 to 60 years old), all of whom had used quantitative descriptive analysis (ISO, 1993) on regular basis over the past 2 years. The panel was calibrated in the use of the chosen attributes during five training sessions. During these sessions, panellists discussed and agreed upon the definitions and how to qualify the

attributes on the scale using commercial soft cheese samples and following the recommendations of the International Dairy Federation (IDF Standard, 1997). Different texture, taste and flavour descriptors, on a 10 cm unstructured line scale anchored with appropriate terms at the left and right ends. For texture attributes, the anchor points were: 1= “almost nothing”, 9= “a lot” and for taste and flavour properties: 1=“barely”, 9=“extremely” perceptible. Test samples of 30 g were presented in plates to the panellist in a randomized order, at 10 °C after 20 days of storage. During tasting, each panel member marked in such scale the perceived intensity of every attribute. Afterwards, the intensities of each descriptor were measured in each scale, in order to assign a numeric value for statistical analysis (Pavón *et al.* 2014).

Statistical analysis

All determinations were performed at least in duplicate. ANOVA test was performed in all experimental determinations and the significance of the results was analyzed by the least significant difference (LSD) test. Differences of $p < 0.05$ were considered to be significant.

RESULTS AND DISCUSSION

Quantification of cholesterol

Cholesterol content of the CB was 10 ± 2 mg mL⁻¹ and 0.7 ± 0.1 mg mL⁻¹ before and after β -CD treatment, respectively. Therefore, the average %ExtChol was 93 ± 1 %. This value agrees with the one previously reported by Kwak *et al.* (2001) for milk cholesterol extraction. These authors report the importance of the homogenisation step in the cholesterol extraction protocol since about 80 % of cholesterol exists in the fat globule. Thus the surface area of the fat globule should be increased for an effective adsorption of β -CD to interact with the cholesterol molecule (Kwak *et al.* 2001).

Chemical cheese analysis and cheese yield determination

Zinc content determination

Zn²⁺ concentrations, % Yield, % Zn²⁺ Fortification and moisture found in cheese samples are shown in Table 1.

As expected, notably higher Zn²⁺ levels ranging from 129±8 to 131±8 mg kg⁻¹, were found in fortified cheese samples (ChZnCl₂ and ChZnSO₄). In addition, Zn²⁺ content values found in fortified cheese and in non-fortified ChWT and ChC-R are comparable with those reported by Aquilanti *et al.* (2012) for Zn²⁺ fortified Squacquerone cheese and Caciotta cheese (Italian cheeses that are similar in manufacture and texture to Cuartirolo soft cheese).

The high % Zn²⁺ fortification found, for both salts used, compared to the low yield of the cheese-making process, is in agreement with the existence of an interaction between Zn²⁺ and casein micelles reported by other authors, who studied the Zn²⁺ binding to caseins (Pomastowski *et al.* 2014). Therefore, most of the added Zn²⁺ remains retained in casein fractions and little is lost in the whey during cheese manufacture.

Assuming a low bioavailability (15 %) of Zn²⁺, a 30 g portion of cheese with the % Zn²⁺ fortification levels (ChZnCl₂ and ChZnSO₄) represents 40% and 28% of the RDA for women and men, respectively. On the other hand, assuming a high bioavailability (50%) of Zn²⁺ it represents about 100% and 90% of the RDA for women and men, respectively. No significant differences were detected between the fortifying agent used and the % fortification value obtained.

The slightly higher yield observed for the cheese obtained from cholesterol-reduced TM compared to ChWT arise as a result of the homogenisation process that involved the cholesterol extraction protocol. Yield increase was previously reported and could be explained by the fact that protein and fat recovery were higher in cheese made from homogenised milk (Kwak *et al.* 2001). Homogenisation creates smaller fat globules with a greater total fat-water interfacial surface area that enabled fat globules to interact with the casein matrix (Everett and Auty 2008).

Moreover, Table 1 shows moisture percentages values for each type of cheese. These values slightly varied among samples where the moisture content of the ChC-R was found to be significantly lower than the rest. Additionally, the presence of the Zn^{2+} salts might promote water retention. Previous reports showed that moisture values of other soft cheese increase with the homogenisation process (Kwak *et al.* 2001, Kwak *et al.* 2002).

Firmness determinations

Cheese samples firmness is shown in Figure 2, in which ChWT firmness was statistically higher compared to the other types of cheese analyzed. These results agree with the observation of a soft and brittle curd during cheese manufacture from β -CD treated CB, also reported by Kwak *et al.* (2002) for cholesterol-reduced Cheddar cheese manufacture. Therefore, these lower firmness values found in cholesterol-reduced cheese samples could be explained by weak coagulum formation as a consequence of CB homogenisation process that caused greater fat globule dispersion in the curd and a reduction in the amount of free casein available to form casein network, resulting in improper curd matting.

On the other hand, no statistical differences in firmness value were found between cheese obtained from cholesterol-reduced TM (ChC-R; ChZnSO₄; ChZnCl₂),

indicating that Zn^{2+} salt fortification does not have an effect on firmness value. Therefore, the Zn^{2+} concentrations added seems not to affect the establishment of rearrangements that lead to increase the coagula firmness.

Digital colorimetric measurement

Figure 3A and 3B exhibit the images of the different cheese samples under study and the colour parameters for each, respectively. In Figure 3B, a decrease in the parameter values a^* and b^* could be observed in cheese samples obtained from the cholesterol-reduced TM in comparison with the values found for ChWT, which reflects a decrease in the intensity of the shades of green (a^*) and yellow colours (b^*). The visual colour differences in ChWT (Figure 3A) are mainly explained by the variation of b^* parameter. Moreover, no differences were found in the values for all analyzed cheese samples for the L^* parameter. These results agree with previous findings by Rudan *et al.* (1998) and Kwak *et al.* (2001). They had reported a change in the cheese structure as a consequence of the reduction of fat particle size by the homogenisation process, which makes the cheese appearance whiter and less yellow than cheese made from unhomogenised milk (Rudan *et al.* 1998, Kwak *et al.* 2001).

Microstructure analysis

The concept of texture from a computational point of view refers to the spatial arrangement of the brightness of the pixels in a region of the image. Therefore, texture parameter study through images reflects changes of intensity values of pixels which might contain information of geometric structure of objects since a great change of intensity values may indicate a change in geometric structure (Zheng *et al.* 2006).

CLSM images of cheese samples sections obtained from the TM with cholesterol-reduced content exhibit a diminution in the values of S, K and $\sigma^2(N)$ and an increment

in the U value, in comparison with the parameter values found for ChWT (Table 2). An increment in the value of U implies a tendency to a uniform distribution of the grey scale of the image, while a diminution of S, K and $\sigma^2(N)$ corresponds to structures in which all the particles are dispersed uniformly over the entire extension of the image (Ingrassia *et al.* 2013). On the other hand, as an observation of images of cheese samples obtained from the TM with cholesterol-reduced content exhibit smaller pores size and pores homogeneously distributed around the entire image (Figure 4), as a result of the process of homogenisation. Nevertheless, CLSM micrographs for ChWT images showed large, irregular and sectorized pores. However, there are no differences in texture parameter values and pore size distributions between the ChC-R, ChZnSO₄ and ChZnCl₂, reflecting the fact that the addition of Zn²⁺ salts has no effect on cheese microstructure, thus concluding that microstructural differences are due exclusively to the homogenisation process.

Meltability

Meltability could be defined as the capability of the cheese to flow or spread upon heating. Figure 5 shows the areas before and after cheese melting process. The results obtained from the difference of areas measured agree with Kwak *et al.* (2001), who found that meltability decreased with homogenisation. In addition, other authors reported that milk homogenisation breaks down fat globule membranes, making the spread of melted fat more difficult (Lelievre *et al.* 1990). Therefore, the mechanical homogenisation step over the CB for cholesterol extraction affects in a negative way the meltability of the resulting cheese. On the other hand, Zn²⁺ salt presence would seem not to affect the melting properties of fortified cheese samples.

Sensory analysis

The sensory attribute of cheese was studied in appearance, texture and flavour; the average data for all sensory descriptors are summarized in Table 3.

Appearance

Cheese appearance was homogeneous with no significant differences among all analyzed cheese samples. Colour was more intense in ChWT than in the other types, which is in consonance with the results found in the digital colour determinations presented. Appearance is one of the most important attributes for cheese and is closely related to consumer's acceptance. From this point of view, whiteness of cholesterol-reduced cheese would grant a desirable feature for the consumer, avoiding the frequent use of chemical strategies, such the addition of titanium dioxide to reach such desirable whiteness (Rudan *et al.* 1998).

Texture

Elasticity for ChZnSO₄ showed a significantly lower value than the rest. Adherence to the palate was higher in the case of fortified cheese than for ChC-R and ChWT, indicating that Zn²⁺ salts increase the resistance to remove the product from the palate. Cohesiveness was lower for ChZnSO₄ and higher for ChC-R. Chewiness and mouthfeel were similar in all samples.

Flavour

Cream odour was detected with more intensity in ChWT than in the other types of cheese. Sweet flavour was detected in low intensity in all the cheese samples studied, ruling out the possibility that a remainder of β -CD could impart sweetness to the cheese mass. Bitter, salty and acid flavours were barely detected in all samples, presenting no significant differences among them. Cream flavour was detected with the same intensity in all samples, indicating that the process of cholesterol extraction does not alter the

organoleptic properties of the traditional cheese. Metallic flavour was not detected and astringency was low for all samples.

Although negative changes in sensory properties of foods fortified with ZnSO_4 have been reported (Salgueiro *et al.* 2002, Boccio and Monteiro 2004), metallic flavour, acid taste and astringency were detected without statistically significant differences among all samples, indicating that fortification with Zn^{2+} salts does not have a significant negative effect on organoleptic characteristics of cheese. In addition, similar sensorial scores were obtained for the fortified cheese samples with the two salts under study, except for elasticity and cohesiveness sensory attributes, in which ChZnCl_2 showed to be more similar to ChWT.

CONCLUSIONS

In the present study, a functional dairy product with a satisfactory level of cholesterol extraction (93%) and a high content of Zn^{2+} level, with adequate physical and sensory characteristics, comparable to those found in traditional Cuartirolo soft cheese, was achieved. However, the differences found between the ChWT and the cholesterol-reduced content cheese samples were due to the mechanical homogenising process used to maximize the cholesterol extraction percentage and were not related with the use of β -CD or with the Zn^{2+} fortification strategy. On the other hand, both Zn^{2+} salts tested showed to be effective and appropriate for Cuartirolo soft cheese fortification. Nevertheless, the use of ZnCl_2 showed to be more adequate from a sensorial point of view.

Based on the overall results we can demonstrate the suitability of the manufacture of a cholesterol-reduced and Zn^{2+} fortified Cuartirolo soft cheese that may meet nutritional needs of risk groups.

ACKNOWLEDGEMENTS

Authors would like to thank those who provide financially and technical support: UNL-CAI+D Program, UNR, Agencia Nacional de Promoción Científica y Tecnológica (PICT-2011-1354), Fundación Nuevo Banco de Santa Fe, Sensory Evaluation Area from Instituto de Tecnología de Alimentos (FIQ-UNL), Lic. Julia Lombardi (UNR-CONICET) and Chemical Engineer Carlos Meinardi (INLAIN-UNL-CONICET). Also we thank to CONICET for the fellowships to Galante Micaela, Lazzaroni Sandra and Pavón Yanina; and the enterprises: Diagramma S.A., Veneto S.A., Ferromet S.R.L., Simes S.A. and Wiener Lab.

The authors are also grateful to the English Area of Facultad de Ciencias Bioquímicas y Farmacéuticas (UNR) for the language correction of the manuscript.

REFERENCES

- ANMAT (2002). *Código Alimentario Argentino. Capítulo VIII: Alimentos lácteos*. Buenos Aires: De la Canal y Asociados.
- Association of Official Analytical Chemists (1990) *Official methods of analysis*, 15th edn. Helrich K, ed. Arlington, VA: Association of Official Analytical Chemists.
- Aquilanti L, Kahraman O, Zannini E, Osimani A, Silvestri G, Ciarrocchi F, Garofalo C, Tekin E and Clementi F (2012) Response of lactic acid bacteria to milk fortification with dietary zinc salts. *International Dairy Journal* **25** 52-59.
- Boccio J and Monteiro J B (2004) Food fortification with iron and zinc: pros and cons from a dietary and nutritional viewpoint. *Revista de Nutrição* **17** 71-78.
- Dias H M A M, Berbic F, Pedrochi F, Baesso M L and Matioli G (2010) Butter cholesterol removal using different complexation methods with beta-

443 cyclodextrin, and the contribution of photoacoustic spectroscopy to the
 444 evaluation of the complex. *Food Research International* **43** 1104-1110.
 445 Doube M, Kłosowski M M, Arganda-Carreras I, Cordelières F P, Dougherty R P,
 446 Jackson J S, Schmid B, Hutchinson J R and Shefelbine S J (2010) BoneJ: Free
 447 and extensible bone image analysis in ImageJ. *Bone* **47** 1076-1079.
 448 El-Din A G, Hassan A, El-Behairy S and Mohamed E (2012) Impact of zinc and iron
 449 salts fortification of buffalo's milk on the dairy product. *World Journal of Dairy*
 450 *& Food Sciences* **7** 21-27.
 451 Everett D W and Auty M A (2008) Cheese structure and current methods of analysis.
 452 *International Dairy Journal* **18** 759-773.
 453 Hess S Y and Brown K H (2009) Impact of zinc fortification on zinc nutrition. *Food &*
 454 *Nutrition Bulletin* **30** 79S-107S.
 455 Ingrassia R, Costa J P, Hidalgo M E, Mancilla Canales M, Castellini H, Riquelme B and
 456 Risso P (2013) Application of a digital image procedure to evaluate
 457 microstructure of caseinate and soy protein acid gels. *LWT - Food Science and*
 458 *Technology* **53** 120-127.
 459 Kahraman O and Ustunol Z (2012) Effect of zinc fortification on Cheddar cheese
 460 quality. *Journal of Dairy Science* **95** 2840-2847.
 461 Kim S Y, Hong E K, Ahn J and Kwak H S (2009) Chemical and sensory properties of
 462 cholesterol-reduced processed cheese spread. *International journal of dairy*
 463 *technology* **62** 348-353.
 464 Kwak H, Jung C, Shim S and Ahn J (2002) Removal of cholesterol from Cheddar
 465 cheese by β -cyclodextrin. *Journal of Agricultural and Food Chemistry* **50**
 466 7293-7298.

467 Kwak H, Nam C and Ahn J (2001) Low cholesterol Mozzarella cheese obtained from
 468 homogenized and β -cyclodextrin-treated milk. *Asian-Australasian Journal of*
 469 *Animal Sciences* **14** 268-275.

470 Lelievre J, Shaker R R and Taylor M W (1990) The role of homogenization in the
 471 manufacture of halloumi and mozzarella cheese from recombined milk.
 472 *International Journal of Dairy Technology* **43** 21-24.

473 Muthukumarappan K, Wang Y C and Gunasekaran S (1999) Modified Schreiber test for
 474 evaluation of Mozzarella cheese meltability. *Journal of Dairy Science* **82** 1068-
 475 1071.

476 Pavón Y L, Lazzaroni S M, Sabbag N G and Rozycki S D (2014) Simultaneous effects
 477 of gelatin and espina corona gum on rheological, physical and sensory properties
 478 of cholesterol-reduced probiotic yoghurts. *International Journal of Food Science*
 479 *& Technology* **49** 2245-4451.

480 Pomastowski P, Sprynskyy M and Buszewski B (2014) The study of zinc ions binding
 481 to casein. *Colloids and Surfaces B: Biointerfaces* **120** 21-27.

482 Rehm C D, Drewnowski A and Monsivais P (2015) Potential Population-Level
 483 Nutritional Impact of Replacing Whole and Reduced-Fat Milk With Low-Fat
 484 and Skim Milk Among US Children Aged 2–19 Years. *Journal of Nutrition*
 485 *Education and Behavior* **47** 61-68.

486 Rudan M A, Barbano D M, Guo M R and Kindstedt P S (1998) Effect of the
 487 modification of fat particle size by homogenization on composition, proteolysis,
 488 functionality, and appearance of reduced fat Mozzarella cheese. *Journal of*
 489 *Dairy Science* **81** 2065-2076.

Salgueiro M J, Zubillaga M, Lysionek A, Caro R, Weill R and Boccio J (2002)
Fortification strategies to combat zinc and iron deficiency. *Nutrition Reviews* **60**
52-58.

Salgueiro M J, Zubillaga M, Lysionek A, Sarabia M I, Caro R, De Paoli T, Hager A,
Weill R and Boccio J (2000) Zinc as an essential micronutrient: a review.
Nutrition Research **20** 737-755.

Soazo M, Pérez L M, Rubiolo A C and Verdini R A (2014) Prefreezing application of
whey protein-based edible coating to maintain quality attributes of strawberries.
International Journal of Food Science & Technology.

WHO/FAO (2005). *Vitamin and mineral requirements in human nutrition*, 230-243.
Switzerland: World Health Organization and Agriculture Organization of the
United Nations.

Zheng C, Sun D-W and Zheng L (2006) Recent applications of image texture for
evaluation of food qualities—a review. *Trends in Food Science & Technology*
17 113-128.

Figure captions

Figure 1. Flow diagrams for Cuartirolo cheese manufacture.

Figure 2. Firmness estimated at 25°C by penetrometry on Cuartirolo cheese samples.

The same letters indicate that there were no significant differences among the samples
($p > 0.05$)

Figure 3. A) Cheese sample images B) Parameters of colour of cheese samples. The
same letters indicate that there was no significant difference among the samples
($p > 0.05$). L* = Lightness (0 = black; 100 = white); a* = red-green component ; b* = yellow-
blue component.

Figure 4. Box plots of the cheese samples pore size distribution. The same letters indicate that there was no significant difference among the samples ($p>0.05$).

Figure 5. Areas before and after melting test. The same letters indicate that there was no significant difference among the samples ($p>0.05$).

Table 1 Data represent the means and standard deviation of cheese chemical composition and cheese yield measurements per sample.*

Table 2 Data represent the means and standard deviation of textural parameters obtained from digital images of different cheese samples: Shannon entropy (S), smoothness (K), uniformity (U), and mean normalized grey-level variance (σ^2 (N)).* All texture parameters are dimensionless.

Table 3 Data represent the means and standard deviation of sensory characteristics of different cheese samples storage at 4°C for 20 days.*

Table 3 Data represent the means and standard deviation of sensory characteristics of different cheese samples storage at 4°C for 20 days.*

Sample	ODOR	COLOR	CHEESE APPEARANCE	ELASTICITY	ADHERENCE	COHESIVENESS
ChWT	4.15±1.70 ^c	4.71±1.75 ^b	8.39±0.55 ^a	4.83±2.12 ^b	2.77±1.37 ^a	5.54±1.77 ^{ab}
ChC-R	2.40±1.96 ^{ab}	2.98±1.35 ^a	7.98±0.98 ^a	5.70±1.47 ^b	3.88±1.58 ^{ab}	6.91±1.19 ^b
ChZnCl ₂	3.54±2.10 ^{cd}	2.71±1.20 ^a	7.23±2.03 ^a	5.17±2.57 ^b	5.01±2.44 ^b	5.65±1.83 ^b
ChZnSO ₄	1.69±1.88 ^a	1.87±1.07 ^a	7.32±2.24 ^a	2.76±2.28 ^a	5.58±3.44 ^b	4.09±1.39 ^a
p-value	0.0190	0.0012	0.4103	0.0243	0.0413	0.0075
Sample	CHEWINESS	MOUTHFEEL	SWEET TASTE	ACID	SALTY	BITER
ChWT	3.34±1.69 ^a	5.43±2.10 ^a	1.39±1.03 ^a	3.37±1.69 ^a	4.29±1.61 ^a	1.98±1.47 ^a
ChC-R	4.29±1.46 ^a	6.39±1.20 ^a	1.51±1.04 ^a	3.10±1.05 ^a	3.67±1.69 ^a	1.98±0.66 ^a
ChZnCl ₂	3.79±0.87 ^a	5.61±1.55 ^a	1.59±1.18 ^a	3.61±2.20 ^a	3.81±1.11 ^a	1.70±1.48 ^a
ChZnSO ₄	2.77±1.01 ^a	6.14±1.72 ^a	1.58±1.11 ^a	3.53±1.62 ^a	3.97±1.94 ^a	2.64±1.40 ^a
p-value	0.1128	0.6060	0.9797	0.9244	0.8717	0.2703
Sample	CREAM FLAVOR	METALLIC FLAVOR	ASTRINGENCY	RESIDUAL FLAVOR		
CWT	3.44±1.84 ^a	2.07±1.67 ^a	1.84±1.93 ^a	4.29±2.87 ^a		
ChC-R	4.38±1.43 ^a	1.34±1.43 ^a	2.30±1.15 ^a	3.49±1.63 ^a		
ChZnCl ₂	4.10±1.92 ^a	1.63±0.88 ^a	2.68±2.16 ^a	4.00±1.70 ^a		
ChZnSO ₄	2.80±1.75 ^a	1.89±1.07 ^a	3.53±1.88 ^a	4.42±2.03 ^a		
p-value	0.1821	0.6292	0.1042	0.8001		

*The same letters indicate that there was no significant difference among the samples (p>0.05)

Table 1 Data represent the means and standard deviation of cheese chemical composition and cheese yield measurements per sample.*

Sample	[Zn ²⁺] (mg kg ⁻¹)	Yield of cheese (% w/w)	Zn ²⁺ Fortification (% w/w)	Moisture (% w/w)
ChWT	(33±2) ^a	(12.2±0.5) ^a	-	(56.4±0.5) ^a
ChC-R	(26±2) ^b	(13.1±0.5) ^{ab}	-	(53.9±0.4) ^b
ChZnCl₂	(129±8) ^c	(13.5±0.1) ^b	(87±7) ^a	(55.8±0.9) ^a
ChZnSO₄	(131±8) ^c	(14±1) ^b	(94±7) ^a	(55.8±0.1) ^a

* The same letters in the same column indicate that there was no significant difference among the samples (p>0.05).

Table 2 Data represent the means and standard deviation of textural parameters obtained from digital images of different cheese samples: Shannon entropy (S), smoothness (K), uniformity (U), and mean normalized grey-level variance (σ^2 (N)).* All texture parameters are dimensionless.

Texture parameters	ChWT	ChC-R	ChZnSO ₄	ChZnCl ₂
S	5.8±0.2 ^a	5.4±0.1 ^b	5.24±0.02 ^b	5.28±0.08 ^b
K	0.043±0.005 ^a	0.017±0.002 ^b	0.014±0.001 ^b	0.016±0.001 ^b
σ^2(N)	2,900±300 ^a	1,100±100 ^b	930±60 ^b	1,050±80 ^b
U	0.023±0.004 ^a	0.031±0.003 ^b	0.031±0.001 ^b	0.030±0.002 ^b

*The same letters indicate that there was no significant difference among the samples (p>0.05)









